





CLASS- 10th

FOUNDATION COURSE

- CBSE, ICSE & OTHER BOARD EXAM
- PRE JEE MAIN & ADVANCED
- PRE MEDICAL (NEET | AIIMS)
- NTSE, KVPY & OTHER COMPETITIVE EXAM





CAREER POINT

Study Material for Pre foundation Class 10

Prepared by Career Point Kota Experts

CONTENTS OF THE PACKAGE AT A GLANCE

Class X

Physics ♦ Electricity ♦ Magnetic effect of Electric Current ♦ Sources of energy ♦ Light-Reflection & Refraction ♦ Human Eye and Colourful world	Chemistry ♦ Chemical Reactions & Equations ♦ Acids, Bases and Salts ♦ Metals & Non-Metals ♦ Carbon and its compounds ♦ Periodic classification of elements	Biology ♦ Nutrition ♦ Respiration ♦ Transportation ♦ Excretion ♦ Control & Coordination ♦ Our Environment ♦ How Do Organisms Reproduce ♦ Heredity and Evolution ♦ Management of Natural Resources
Mathematics [Set-1] ♦ Real Numbers ♦ Polynomials ♦ Pair of Linear Equations in two variables ♦ Quadratic equations ♦ Arithmetic Progressions ♦ Triangles ♦ Coordinate Geometry	Mathematics [Set-2] ♦ Introduction to Trigonometry ♦ Some Applications of Trigonometry ♦ Circles ♦ Constructions ♦ Areas Related to Circles ♦ Surface Areas and Volumes ♦ Statistics ♦ Probability	Mental Ability ♦ logical sequence of words ♦ pyramid test ♦ calendar test ♦ clock test ♦ data sufficiency ♦ data redundancy ♦ cube test ♦ dice test ♦ mirror & water image ♦ paper cutting & folding ♦ figure partition & counting ♦ completion & formation of figures ♦ embedded figure ♦ non-verbal series ♦ non-verbal analogy ♦ non-verbal classification ♦ dot situation

Social Science [Set-1]

History

- ◆ The Rise of Nationalism in Europe
- ◆ The Nationalist Movement in Indo-China
- ◆ Nationalism in India
- ◆ The Making of a Global World
- ◆ The Age of Industrialisation
- ◆ Work, Life & Leisure - Cities in Contemporary World
- ◆ Print Culture and The Modern World
- ◆ Print Culture and The Modern World

Economics

- ◆ Development
- ◆ Sectors of the Indian Economy
- ◆ Money And Credit
- ◆ Globalisation And The Indian Economy
- ◆ Consumer Right

Social Science [Set-2]

Geography

- ◆ Resource & Development
- ◆ Forest And Wildlife Resources
- ◆ Water Resources
- ◆ Agriculture
- ◆ Minerals And Power Resources
- ◆ Manufacturing Industries
- ◆ Lifelines of National Economy

Civics

- ◆ Power Sharing
- ◆ Federalism
- ◆ Democracy & Diversity
- ◆ Gender, Religion and Caste
- ◆ Popular Struggles and Movements
- ◆ Political Parties
- ◆ Outcomes of Democracy
- ◆ Challenges to Democracy

English [Set-1]

- ◆ Letter writing
- ◆ Article writing
- ◆ Story writing
- ◆ Integrated grammar
- ◆ The frog and the nightingale
- ◆ Mirror
- ◆ Not marble nor the gilded monuments
- ◆ Two gentlemen of verona
- ◆ Mrs. Packletide's tiger
- ◆ The letter
- ◆ The dear departed
- ◆ The diary of a young girl (term-1)
- ◆ The story of my life (term-1)
- ◆ Ozymandias
- ◆ The rime of ancient mariner
- ◆ Snake
- ◆ Shady plot
- ◆ Patol babu
- ◆ Virtually true
- ◆ Julius caesar
- ◆ The diary of a young girl (term-2)
- ◆ The story of my life (term-2)
- ◆ Figures of speech

English [Set-2]

- ◆ Noun
- ◆ Pronoun
- ◆ Adjective
- ◆ Adverb
- ◆ Determiners
- ◆ Connectors
- ◆ Prepositions
- ◆ Tense
- ◆ Passivization
- ◆ Reported speech
- ◆ Modals
- ◆ Subject verb concord
- ◆ Conditionals
- ◆ Error spotting
- ◆ Reading comprehension
- ◆ Fillers/sentence completion
- ◆ Cloze test
- ◆ Missing sen./jumbled sentences & paragraphs
- ◆ Sentence improvement
- ◆ Synonyms
- ◆ Antonyms
- ◆ Idioms and phrases
- ◆ Phrasal verbs
- ◆ One word substitution

Note to the Students



Career Point offers this must have Study Package in Physics, Chemistry and Mathematics to meet the complete curriculum needs of engineering aspirants. The set comprises of 18 books: **Physics** - set of 3 books for class 11 and set of 3 books for Class 12; **Chemistry** - set of 3 books for class 11 and set of 3 books for Class 12 and **Mathematics** - set of 3 books for class 11 and set of 3 books for Class 12. The set caters to the different requirements of students in classes XI and XII. It offers complete and systematic coverage of **JEE Main** and **JEE Advanced** syllabi and aims to provide firm foundation in learning and develop competitive edge in preparation of the JEE and other engineering entrance examinations.

COMPONENTS OF EACH CHAPTER

These books are designed with an engaging and preparation-focused pedagogy and offer a perfect balance of conceptual learning and problem solving skills.


Mind Map

Each chapter contains many articles (Concepts, Theories etc.). Mind map interconnect all these articles logically. By this student can understand whole chapter articles interconnectivity clearly in a single picture frame.



Theory & Concepts

Each chapter consist of exhaustive theory which gives conceptual clarity and command over topics. Appropriate explanation of theory with the help of images, diagrams, flowcharts, mind maps, info graphics, and tables.



MAGNETIC EFFECT OF ELECTRIC CURRENT

Magnet & Magnetism

◆ Magnet


The substance which attract Ni, Fe, Co, etc and align in geographical north south direction when suspended freely.

Two bodies even after being neutral (showing no electric interaction) may attract / repel strongly if they have a special property. This property is known as magnetism. This force is called magnetic force. Those bodies are called magnets.

• Origin of the Magnet

Around 120 A.D. near Magnesia in Asia Minor, an ore of iron Fe_3O_4 was found in abundance. This ore of iron, which was found attracting small pieces of iron, was called Magnetite. Hence, the name 'magnet' was given to the pieces that exhibited this property.

• Types of Magnet



Competitive Level

Competitive level is specially designed for competition exam requirements and to better understanding the concepts, well explained theory, clearly explained formulas with good number of quality examples are given in this.

COMPETITIVE LEVEL

- **Magnitude of magnetic field produced by a straight current-carrying conductor:** The magnitude of magnetic field (or strength of magnetic field) B produced by an infinitely long conductor in vacuum at a distance r from it, is given by:

$$B = \frac{\mu_0 I}{2\pi r}$$

where, B = Magnetic field strength

μ_0 = Permeability of vacuum (a constant)

I = Current (flowing in conductor) and

r = Distance from the conductor (where magnetic field is measured).

The SI unit of magnetic field B is Tesla which is denoted by the symbol T (1 tesla is equal to 1 newton per ampere per metre).

The CGS unit of magnetic field B is Gauss which is denoted by the symbol G. (1 Tesla = 10^4 Gauss)

Permeability of vacuum μ_0 is $4\pi \times 10^{-7}$ tesla metre per ampere.

In Chapter Example

To understand the application of concepts, there is *in chapter solved example* are given. It contains large variety of all types of solved examples with explanation to ensure understanding the application of concepts.

Ex.31 Compute the heat generated while transferring 96000 coulomb of charge in one hour through a potential difference of 50 V.

Sol. The amount of heat (H) produced is given by the Joule's law of heating as $H = VIt$

where, Voltage, $V = 50$ V

Time, $t = 1 \text{ h} = 1 \times 60 \times 60 \text{ s} = 3600 \text{ s}$

Amount of current, $I = q/t = 96000 / 3600 = (80/3) \text{ A}$

$$H = 50 \times \frac{80}{3} \times 60 \times 60 = 4.8 \times 10^6 \text{ J}$$

Therefore, the heat generated is $4.8 \times 10^6 \text{ J}$.

Ex.32 An electric iron of resistance 20Ω takes a current of 5 A. Calculate the heat developed in 30 s.

Sol. The amount of heat (H) produced is given by the Joule's law of heating as $H = VIt$

where, Current, $I = 5 \text{ A}$

Practice Exercises

Includes three sets of exercises covering all the topics. Helps the students to assess their strengths and weaknesses and work on them accordingly. Separate exercises for subjective as well as objective questions and previous year competitive exams questions (NTSE, Olympiads)

EXERCISE-1

➤ Very Short Answer Type Questions

- Q.1** What is represented by joule/coulomb?
- Q.2** What is the function of electric switch in electric circuit?
- Q.3** A wire of resistivity 10 ohm meter is stretched to double its length. What is its new resistivity?
- Q.4** What is the SI unit of Power?
- Q.5** A charge of 2C moves between two plates, maintained at a p.d of 1V. What is the energy acquired by the charge?

➤ Short Answer Type Questions – Type I

- Q.6** Draw a schematic diagram of a typical electric circuit comprising a cell, an electric bulb, an ammeter and a plug key.
- Q.7** What makes the electric charge to flow in a conducting metallic wire?
- Q.8** Two wires of same material and same

Q.13 A TV set shoots out a beam of electrons. The beam current is $10\mu\text{A}$.

- (a) How many electrons strike the TV screen in each second?
- (b) How much charge strikes the screen in a minute?

Q.14 (a) Why do copper or aluminium wires generally used for electrical transmission and distribution purposes?

- (b) Two wires, one of copper and other of manganin, have equal lengths and equal resistances. Which wire is thicker? Given that resistivity of copper is lower than that of manganin.

Q.15 Two conductors X and Y of circular cross-section have radii in the ratio of 1: 2, length in the ratio 1: 3 and resistivities in the ratio of 3: 1, then in which ratio their resistances are?

Q.16 Given n resistors each of resistance r . How will you combine them to get the

(i) Maximum and

(ii) Minimum effective resistance? What is the ratio of the maximum to minimum resistance?


EXERCISE-2

- Q.1 When a body is negatively charged by friction, it means
 (A) the body has acquired excess of electrons
 (B) the body has acquired excess of protons
 (C) the body has lost some electrons
 (D) the body has lost some neutrons
- Q.2 When the distance between the charged particles is halved, the force between them becomes
 (A) One-fourth (B) Half
 (C) Double (D) Four times
- Q.8 Deuteron and α -particle are put $1\text{ \AA}$ apart in air. Magnitude of intensity of electric field due to deuteron at α -particle is
 (A) Zero
 (B) 2.88×10^{11} newton/coulomb
 (C) 1.44×10^{11} newton/coulomb
 (D) 1.44×10^{11} newton/coulomb
- Q.9 What is not true for electric charge :
 (A) Electric charge is scalar quantity
 (B) Charge on a body may be +ve or -ve
 (C) S.I. unit of charge is coulomb
 (D) One coulomb is charge of one electron

EXERCISE-3

- Q.1 In the given circuit diagram, the value of resistance X in ohm when the bridge is balanced will be

[Raj./ NTSE Stage-I/2005]



- (A) 4 (B) 8
 (C) 10 (D) 12

- (A) high melting point and high specific resistance
 (B) high melting point and low specific resistance
 (C) low melting point and high specific resistance
 (D) low melting point and high specific resistance

- Q.5 A uniform wire when connected directly across a 220V line produces heat H per second. If the wire is divided into n-parts and all parts are connected in parallel across a 220 V line, the heat produced per second will be

[Haryana/NTSE Stage-I/2014]

- (A) H/n (B) H/n^2
 (C) n^2H (D) nH

Answer key

Answer key is provided at the end of the exercise sheets.

ANSWER KEY

EXERCISE - 1

3. 40ω
 5. 2 J
 8. $\frac{R_1}{R_2} = \left(\frac{r_2}{r_1}\right)^2$
 10. $1.44 \times 10^6\text{ J}$
 12. $2qv/\ell$
 13. (a) 6.25×10^{13} (b) $6 \times 10^{-4}\text{C}$

EXERCISE - 2

Ques.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Ans.	A	D	A	D	C	C	C	C	D	A	A	A	C	A	B
Ques.	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
Ans.	D	C	D	A	A	B	D	D	B	C	C	A	C	C	A


EXERCISE - 3

Ques.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Ans.	B	A	B	B	B	C	A	C	A	A	C	B	D	B	A
Ques.	16	17	18	19	20	21	22	23	24						
Ans.	B	A	A	B	A	B	D	A	C						

ELECTRICITY


Chapter Outline

- ❖ Electric Charge
- ❖ Static & Current Electricity
- ❖ Electric field
- ❖ Electric Potential and Potential Difference
- ❖ Electric Potential Energy
- ❖ Electric Current & Circuits
- ❖ Ohm's Law
- ❖ Resistance of a Conductor
- ❖ Combination of Resistances
- ❖ Heating Effect of Electric Current



MIND MAP

ELECTRICITY



ELECTRICITY

Electric Charge

◆ Definition

Electric charge may be defined as the intrinsic property of certain fundamental particles (electron, proton, etc.) due to which they produce electric and magnetic effects.

◆ Charge on a Macrobody

Excess or deficiency of electrons in a body is equal to the charge on a macrobody. A body having excess of electrons is negatively charged and a body having deficiency of electrons is positively charged.

From the study of atomic structure, we know that an atom consists of a central part called nucleus and around the nucleus there are a number of electrons revolving in different paths or orbits. The nucleus contains protons and neutrons. A proton is a positively charged particle while a neutron has no charge. Therefore, the nucleus of an atom bears a positive charge. An electron is a negatively charged particle having negative charge equal to the positive charge on a proton. Normally, the number of electrons is equal to the number of protons in an atom. Therefore, an atom is neutral as a whole; the negative charge on electrons cancelling the positive charge on protons. This leads to the conclusion that under ordinary conditions, a body is neutral i.e. it exhibits no charge.

When this equity or balance is disturbed by removing or supplying electrons, the body acquires a net charge. The body will acquire a positive or negative charge depending upon whether electrons are removed from it or added to it.

◆ Types of Electric Charge

There are two types of charges:

- (i) Positive charge - A body having deficiency of electrons as compared to proton.
- (ii) Negative charge- A body having excess of electrons as compared to proton.

◆ Charging a body

There are a number of methods to charge a body as:

- Charging by friction
- Charging by conduction
- Charging by induction, etc.

◆ We will discuss charging by friction


Whenever two bodies (at least one non conductor) are rubbed against each other, heat is produced due to friction present between them. Due to this heat produced, electrons in both the bodies are excited. The body having more electron affinity attracts some of the electrons from other body. Both the bodies develop equal and opposite charges by this method.

POSITIVE CHARGE	NEGATIVE CHARGE
1. Glass Rod	1. Silk cloth
2. Fur or woolen cloth	2. Ebonite, Amber, Rubber rod
3. Woolen coat	3. Plastic seat
4. Woolen carpet	4. Rubber shoes
5. Nylon or Acetate	5. Cloth
6. Dry hair	6. Comb

Note: The object in above table must be in given pair.

◆ Properties of Electric Charge

- Like charges repel and unlike charges attract each other.
- Charge is a scalar quantity
- Charge is always quantized: The amount of charge on a charged body is always in integral multiple of the elementary charge and the fractional multiple is not possible.
- Charge is conserved.
- Charge is always associated with mass.
- Total charge of system remains conserved.



◆ Unit of Charge

The S.I. unit of charge is coulomb abbreviated as C. One coulomb of charge is equal to the charge on 625×10^{16} electrons.

1 coulomb = charge on 625×10^{16} electrons

or 6.25×10^{18} electrons

Thus, when we say that a body has a positive charge of one coulomb (i.e. + 1C) it means that the body has a deficit of 625×10^{16} electrons from the normal due to share.

Note: The attraction and repulsion of electric charges can be used in experiments to demonstrate the existence of two types of charge. Note that it is possible for a charged object to attract an uncharged object that is free to move but it cannot repel an uncharged object because it is neutral.

Ex.1 Calculate the number of electrons constituting one coulomb of charge.

Sol. Number of electrons constituting one coulomb of charge:-

From $q = ne$

$$n = \frac{q}{e} = \frac{1\text{C}}{1.6 \times 10^{-19}\text{C}} = 6.25 \times 10^{18} \text{ electrons}$$

Static and Current Electricity

◆ Static electricity

A branch of physics which deals with the study of the electric charges at rest and their effects is known as electrostatic or static electricity.


◆ Current electricity

A branch of physics which deals with the study of the electric charges in motion and their effects is known as current electricity.

COMPETITIVE LEVEL

Coulomb's Law

Charles Augustine De Coulomb studied the interaction forces of charged particles in detail in 1784. He used a torsion balance. On the basis of his experiments he established Coulomb's law. According to this law, the force of attraction or repulsion between two stationary point charges is directly proportional to the product of charges and inversely proportional to the square of distance between them. This force acts along the line joining the two. If q_1 & q_2 are charges in consideration, r , the distance between them and F , the force acting between them



Then, $F \propto q_1 q_2$

$F \propto 1/r^2$

$$\therefore F \propto \frac{q_1 q_2}{r^2}$$

$\Rightarrow F = k \frac{q_1 q_2}{r^2}$, where $k = \text{constant}$.

$$K = \frac{1}{4\pi\epsilon_0\epsilon_r} = \frac{9 \times 10^9}{\epsilon_r} \text{ Nm}^2\text{C}^{-2}$$

Where,

ϵ_0 = Electric permittivity of vacuum or air

$$= 8.85 \times 10^{-12} \text{ C}^2\text{N}^{-1} \text{ m}^{-2} \text{ and}$$


K or ϵ_r = Relative permittivity or Dielectric constant

or Specific inductive capacity

$$\epsilon_r = \frac{\epsilon}{\epsilon_0} \Rightarrow \epsilon = \epsilon_0 \epsilon_r$$

[Newton's law for particles is analogous to coulomb's law for rest charges. The difference is that Newton's law gives attraction force while coulomb's law gives attraction as well as repulsion force]

Direction: Direction of the force acting between two charges depends upon their nature and it is along the line joining two charges.



$$\vec{F}_{21} = \text{force on } q_2 \text{ due to } q_1 \quad \vec{F}_{21} = \frac{q_1 q_2}{4\pi\epsilon_0\epsilon_r r_{12}^2} \hat{r}_{12} \quad \dots\dots(A)$$

(where $\hat{r}_{12}$ is a unit vector pointing from q_1 to q_2)

$\vec{F}_{12}$ = Force on q_1 due to q_2


$$\vec{F}_{12} = \frac{q_1 q_2}{4\pi\epsilon_0\epsilon_r r_{12}^2} \hat{r}_{21} \quad \dots\dots(B)$$

(where $\hat{r}_{21}$ is a unit vector pointing from q_2 to q_1)


$\Rightarrow$ Electric force between two charges does not depend on neighboring charges.

⇒ When two charges (Q_1, Q_2) are placed some distance apart. Neutral point is nearer to smaller charge, in between Q_1 and Q_2 if charges are like and away from charge if charges are unlike.

⇒ System of following charges is in equilibrium, if



$$Q_1 = -\frac{Q}{4}$$



$$Q_2 = -\frac{Q}{\sqrt{3}}$$


⇒ For a three charge system it is not possible that all charges are in **stable** equilibrium.

⇒ Coulomb's law is similar to Newton's gravitation law.

⇒ If $F_g = F_e$ for two identical charges then,

$$\frac{q}{m} = \sqrt{4\pi\epsilon_0 G}$$

Ex.2 Ten charged particles are kept fixed on the X axis at point $x = 10$ mm, 20 mm, 30 mm, 100 mm. The first particle has a charge 10^{-8} C, the second 8×10^{-8} C, the third 27×10^{-8} C and so on. The tenth particle has a charge 1000×10^{-8} C. Find the magnitude of electric force acting on a 1 C charge placed at the origin.



Sol. Force of $1C$ charge = $\frac{Kq_1 \times 1}{(10 \times 10^{-3})^2} + \frac{Kq_2 \times 1}{(20 \times 10^{-3})^2} + \frac{Kq_3 \times 1}{(30 \times 10^{-3})^2} + \dots$

$$= \frac{K \times 10^{-8}}{10^{-4}} \left[\frac{1^3}{1^2} + \frac{2^3}{2^2} + \frac{3^3}{3^2} + \dots + \frac{10^3}{10^2} \right] = 9 \times 10^9 \times 10^{-4} \times 55 = 4.95 \times 10^7 \text{ N.}$$

◆ Electric Field

Electric field due to a given charge is defined as the space around the charge in which electrostatic force of attraction or repulsion due to charge can be experienced by any other charge. If a test charge experiences no force at a point, the electric field at that point must be zero.

◆ Electric Field Intensity

Electric field intensity at any point is the strength of electric field at that point. It is defined as the force experienced by unit positive charge placed at that point.

If F is the force acting on a test charge $+q_0$ at any point then electric field intensity at this point is given by

$$E = \frac{F}{q_0}$$

Electric field intensity is a vector quantity and its S.I. unit is Newton per coulomb (N/C).

◆ Electric Field Strength due to Point Charge

As discussed earlier, if we find electric field due to a point charge at a distance x from it. Its magnitude can be given as

$$E = \frac{kq}{x^2}$$

Ex.3 What is the magnitude of a point charge due to which the electric field 30 cm away has the magnitude 2 *Newton/Coulomb* [$1/4 \pi \epsilon_0 = 9 \times 10^9 \text{ Nm}^2 \text{ C}^{-2}$]

Sol. By using $E = \frac{1}{4\pi\epsilon_0} \cdot \frac{Q}{r^2}$; $2 = 9 \times 10^9 \times \frac{Q}{(30 \times 10^{-2})^2} \Rightarrow Q = 2 \times 10^{-11} \text{ C}$

Ex.4 A particle of mass $9 \times 10^{-31} \text{ kg}$ and a negative charge of $1.6 \times 10^{-19} \text{ coulomb}$ projected horizontally with a velocity of 10^5 m/s into a region between two infinite horizontal parallel plates of metal. The distance between the plates is 0.3 cm and the particle enters 0.1 cm below the top plate. The top and bottom plates are connected respectively to the positive and negative terminals of a 30 volt battery. Find the component of velocity of the particle just before it hits one of the plates. (Given $E = V/d$)

Sol. We know that between two parallel plates electric field can be given as

$$E = \frac{V}{d}$$

Here $V = 30 \text{ volt}$ and $d = 0.3 \text{ cm} = 3 \times 10^{-3} \text{ m}$

Thus, we have $E = \frac{30}{3 \times 10^{-3}} = 10^4 \text{ N/C}$

Force on the particle of negative charge moving between the plates

$$F = e \times E = 1.6 \times 10^{-19} \times 10^4 = 1.6 \times 10^{-15} \text{ N.}$$

The direction of force will be towards the positive plate i.e., upward.

Now, acceleration of the particle is

$$a = \frac{eE}{m}$$

$$\text{or } a = (1.6 \times 10^{-15}) / (9 \times 10^{-31})$$

$$\text{or } a = 1.77 \times 10^{15} \text{ m/sec}^2$$

As the electric field intensity E is acting in the vertical direction, the horizontal velocity v of the particle remains same. If y is the displacement of the particle in upward direction, we have

$$y = \frac{1}{2} at^2$$

Here, $y = 0.1 \text{ cm} = 10^{-3} \text{ m}$, $a = 1.77 \times 10^{15} \text{ m/sec}^2$

$$\text{Thus } 10^{-3} = \frac{1}{2} \times (1.77 \times 10^{15}) (t^2)$$

Solving, we get $t = 1.063 \times 10^{-9}$ second

Component of velocity in the direction of field is given by,


$$v_y = at$$

$$= (1.77 \times 10^{15}) (1.063 \times 10^{-10})$$

$$= 1.881 \times 10^6 \text{ m/s.}$$

Electric Potential and Potential Difference

The flow of electricity in a circuit can be regarded very much similar to the flow of water in a pipe. The water pipe is analogous to the electric conductor, while the amount of water flowing through a given point per second corresponds to electric current. Figure below shows how the pump (P) builds up and maintains pressure by lifting water from a tank (B) to the reservoir (A) through the pipe (R). Note that along the pipe, different points are at different pressure. Water in the pipe flows from, say point C to point D, only when the pressure at C is greater than that at D. Thus, when the valve (V) is open, water would start flowing into the reservoir.



In the same manner electrons will move along a wire only if there is a difference of electric pressure called potential difference along the conductor. This difference of potential is produced by the cell or a battery, which acts like a water pump in the circuit.

The chemical action within the cell generates the difference in potential between the electrodes, which sets the electrons in motion and produces the current.

Electric Potential: Work done in bringing a unit positive charge from infinity to any point in an electric field of another charge is termed as potential at that point i.e. if,

$$W = \text{work done in bringing a positive charge } q_0 \text{ from infinity to that point, then, } V = \frac{W}{q_0}$$

Since work is measured in joule and charge in coulomb, therefore electric potential is measured in joule per coulomb (J/C).

Joule per coulomb occurs so often in our study of electricity, so it has been named as volt, in honour of the scientist Alessandro Volta (the inventor of the voltaic cell).

$$1 \text{ Volt} = \frac{1 \text{ joule}}{1 \text{ coulomb}}$$

Potential is a scalar quantity, therefore it is added algebraically. For a positively charged body potential is positive and for a negatively charged body potential is negative.

- We can say potential is the electrical state of a conductor which determines the direction of flow of charge when the two conductors are kept in contact.

Some Important Points

- (i) Electric potential at infinity is taken to be zero.
- (ii) It is not path dependent quantity, it simply depends upon the starting and end points.
- (iii) It is a scalar quantity.
- (iv) Unit: Volt or Joule/Coulomb
- (v) Dimension: $[M^1 L^2 T^{-3} A^{-1}]$
- (vi) Potential due to a positive charge is positive and potential due to a negative charge is negative. Here, potential being positive and negative implies whether work is done on the charge or done by the charge respectively.
- (vii) Potential due to a point charge Q at a distance r is

$$V = \frac{1}{4\pi\epsilon_0} \frac{q}{r} \Rightarrow V \propto \frac{1}{r}$$

- (viii) Total potential at a point due to a group of charges is scalar sum of individual potentials

$$V_p = V_1 + V_2 + \dots V_n$$

- (ix) Electric field is gradient of electric potential at that point.

$$E = - \frac{dv}{dr}$$

Note: The negative sign implies that direction of electric field is in the direction of decreasing potential.

- (x) Potential of earth is taken to be zero.

Potential difference: The work done in taking a charge from one point to the other in an electric field is called the potential difference between two points.


Thus, if W be work done in moving a charge q_0 from B to A then the potential difference is given by-

$$V_A - V_B = \frac{W}{q_0}$$

Note: Do not confuse e.m.f. and potential difference (V), even though they have the same units. E.m.f., $\mathcal{E}$, is provided by a source of electrical energy, but potential difference (V) refers to the electrical energy converted to other forms by a circuit component.

Electric Potential Energy

Consider a charge Q placed at a point P as shown in figure. If another charge q of the same sign is now brought from a very far away distance (infinity) to point O near P, then charge q will experience a force of repulsion due to charge Q . If charge q is still pushed towards P, work is done. This work done is the potential energy of the system of these two charges.



Thus, the electric potential energy of a system of charges is defined as the amount of work done in bringing the various charges from infinite separation to their present positions to form the required system. It is denoted by U . For the system of two charges separated by distance r as shown in figure, the electric potential energy is given by:

$$U = \frac{kQq}{r}$$

Electric potential energy is the form of energy, therefore, it is measured in joule (J). The potential difference is measured by means of an instrument called voltmeter. The voltmeter is connected in parallel across the points where the potential difference is to be measured. A voltmeter has a high resistance so that it takes a negligible current from the circuit.

Work done in bringing a charge Q from infinity to that point is,

$$W = QV$$

Where, V is potential at that point.

Ex.5 How much energy is given to each coulomb of charge passing through a 6 V battery?

Sol. Energy required to each coulomb of charge passing through a 6 V battery is $U = VQ = 6 \text{ V} \times 1 \text{ C} = 6 \text{ J}$.

Ex.6 What is meant by saying that the potential difference between two points is 1 V?

Sol. If 1 Joule of work is required to move charge of 1C from one point to another, then it is said that the potential difference between the two points is 1V.


Ex.7 Name a device that helps to maintain a potential difference across a conductor. How it helps to maintain the current in a given electric circuit? What energy changes takes place in it?

Sol. To maintain a potential difference across a conductor we use a device known as cell. The chemical action within a cell generates the potential difference across the terminals of the cell, even when no current is drawn from it. When the cell is connected to a conducting circuit element, the potential difference sets the charges in motion in the conductor and produces an electric current. In order to maintain the current in a given electric circuit, the cell has to expend its chemical energy stored in it hence cell converts chemical energy into electrical energy to move charges in electric circuit.

Electric Current & Circuits

◆ Charges in motion

When two charged bodies at different potentials are connected by a conducting wire, electrons flow from the body at lower potential to the one at higher potential till the potentials of both the bodies become equal i.e., potential difference between the two bodies becomes zero. Figure below, shows the two oppositely charged conductors A and B which are held on stands. The positively charged conductor A is said to be at higher potential and the negatively charged conductor B is said to be at lower potential.



It means that there exists a potential difference between the conductors A and B. Therefore, on joining positively charged conductor A to negatively charged conductor B, negative charge starts flowing from conductor B to conductor A. Flow of charge will stop when both the conductors A and B acquire the same potential.

This rate of flow electric charge from one body to another through a conductor such as metal wire is called electric current and its direction is opposite to direction of flow of electrons.

Or

The quantity of charge passing through a given point of the conductor in one second is called electric current.

Thus, if Q is the charge which flows through a conductor in time t , then the electric current is given by

$$\text{Current (I)} = \frac{\text{Charge(Q)}}{\text{Time(t)}}$$

or
$$I = \frac{(Q)}{(t)}$$

or
$$Q = It$$

Note: The electric current is a scalar quantity.

◆ Unit of Current

S.I. unit of current is ampere, which is denoted by letter A.

Current is said to be one ampere if one coulomb of charge flows through any cross section of a conductor in one second, i.e.

$$1 \text{ ampere} = \frac{1 \text{ coulomb}}{1 \text{ second}} \quad \text{or} \quad 1 \text{ A} = 1 \text{ C s}^{-1}$$

◆ Smaller Units of Current

$$1 \text{ milliampere (mA)} = 10^{-3} \text{ A}$$


$$1 \text{ microampere (A)} = 10^{-6} \text{ A}$$

◆ Direction of Electric Current


When electricity was invented a long time back, it was known that there are two types of charges: positive charges and negative charges, but the electron had not been discovered at that time. So, electric current was considered to be a flow of positive charges and the direction of flow of the positive charges was taken to be the direction of electric current. Thus, the conventional direction of electric current is from positive terminal of a cell (or battery) to the negative terminal through the circuit.

◆ How the Current Flows in a Wire

As electric current is the flow of electrons in a metal wire (or conductor) when a cell or battery is connected across its ends. A metal wire has plenty of free electrons in it. When the metal wire has not been connected to a source of electricity like a cell or a battery, then the electrons present in it move at random in all the directions between the atoms of the metal wire as shown in figure below.



When a source of electricity like a cell or a battery is connected between the ends of the metal wire, then an electric force acts on the electrons present in the wire. Since the electrons are negatively charged, they start moving from negative end to the positive end of the wire and this flow of electrons constitutes the electric current in the wire.




Note: Electric eel fish can generate around 600V and 1 ampere of current through the water it uses this capacity to electrocute and kill its prey.

◆ How to get a Continuous Flow of Electric Current

It is due to the potential difference between two points that an electric current flows between them. The simplest way to maintain a potential difference between the two ends of a conductor so as to get a continuous flow of current is to connect the conductor between the terminals of a cell or a battery. Due to the chemical reactions going on inside the cell or battery, a potential difference is maintained between its terminals and this potential difference drives the current in a circuit.

Ex.8 If 'q' coulombs of charge travel through a conductor of length ' ℓ ' m with a velocity of ' v ' m/s, what is the current flowing through the conductor?

Sol. Charge passing through the conductor = 'q' coulomb



We know that current, $i = \frac{q}{t}$ (i)

and speed, $v = \frac{\ell}{t} \Rightarrow t = \frac{\ell}{v}$

Substituting the value of 't' in equation (i)

We get $i = \frac{q}{t} = \frac{q}{\left(\frac{\ell}{v}\right)} = \frac{q \times v}{\ell}$

$\therefore$ The current flowing through conductor = qv/ℓ

Ex.9 An electron of charge 'e' C moves in a circular orbit at a frequency of 'n' revolution per second. Then find the strength of the current.

Sol. Let P be a reference point in the orbit

$$i = \text{rate of flow charge} = \frac{q}{t}$$

Consider, $t = 1\text{ s}$



In 1 second the electron completes 'n' revolutions (frequency is 'n' rev/s)


$\Rightarrow$ In 1 second the electron passes through 'P' for 'n' times.

$\therefore$ charge through 'P' in 1 second (q) = ne, where 'e' is charge of electron.

$$\therefore i = \frac{q}{t} = \frac{ne}{1} = ne \text{ (amp.)}$$


Electric circuit

A continuous path consisting of conducting wires and other components (like lamps, bulbs etc.) between the terminals of a battery, along which an electric current flows, is called a circuit.




◆ Open Electric Circuit

An electric circuit through which no electric current flows is known as open electric circuit. The electric circuit will be open circuit if the plug of the key is taken out or if the connecting wires break from any point.



◆ Closed Circuit

An electric circuit through which electric current flows continuously is known as closed circuit.



COMPETITIVE LEVEL**◆ Dependency of a Resistance on Temperature**

If R_0 is the resistance of the conductor at 0°C and R_t is the resistance of the conductor at $t^\circ\text{C}$ then the relation between R_0 and R_t is given by,

$$R_t = R_0(1 + \alpha\Delta t) \quad [\text{Here } \Delta t = t - 0 = t]$$

$$\text{or } \alpha = \frac{R_t - R_0}{R_0 t}$$

Here, α = Temperature Coefficient of Resistance, (SI unit = $^\circ\text{C}^{-1}$)

t = temperature in $^\circ\text{C}$

Ex.15 Temperature coefficient of resistance is 0.00125 per $^\circ\text{C}$. Resistance at 300K is 1Ω . What will be the temperature in (K) when resistance becomes 2Ω ?

$$\text{Sol. } \alpha = \frac{R_2 - R_1}{R_1 t_2 - R_2 t_1}$$

$$\Rightarrow 0.00125 = \frac{2 - 1}{1 \times t_2 - 2 \times 27} \quad [t_1 = 300 \text{ K} = 27^\circ\text{C}]$$

$$\Rightarrow t_2 = 854^\circ\text{C}$$

$$t_2 = 1127 \text{ K}$$

◆ Resistivity

$$\rho = \frac{R \times A}{L}$$

By using this formula, we will now obtain the definition of resistivity. Let us take a conductor having a unit area of cross-section of 1 m^2 and a unit length of 1 m . So, putting $A = 1$ and $L = 1$ in above equation we get:

Resistivity, $\rho = R$

- **Definition of resistivity:** The resistivity of a substance is numerically equal to the resistance of a rod of that substance which is 1 metre long and 1 metre square in cross-section.

Unit of resistivity,

$$[\rho] = \frac{\text{ohm} \times (\text{metre})^2}{\text{metre}} = \text{ohm} \cdot \text{metre}$$

The S.I. unit of resistivity is ohm-metre which is written in symbols as $\Omega\text{-m}$. Resistivity of a substance does not depend on its length or thickness. It depends only on the nature of the substance. The resistivity of a substance is its characteristic property. So, we can use the resistivity to compare the resistances of two or more substances.

- **Importance of resistivity:** A good conductor of electricity should have a low resistivity and a poor conductor of electricity should have a high resistivity. The resistivity of alloy are much more higher than those of the pure metals.

It is due to their high resistivities that manganin and constantan alloys are used to make resistance wires used in electronic appliances to reduce the current in an electrical circuit.

Nichrome alloy is used for making the heating elements of electrical appliances like electric irons, room-heaters, water-heaters and toasters etc. because it has very high resistivity and it does not undergo oxidation (or burn) even when red-hot.

COMPETITIVE LEVEL

- **Effect of temperature on resistivity:** The resistivity of conductors (like metals) is very low. The resistivity of most of the metals increases with temperature. On the other hand, the resistivity of semi-conductors like silicon and germanium is in between those of conductors and insulators and decreases on increasing the temperature. Semi-conductors are proving to be of great practical importance because of their marked change in conducting properties with temperature and impurity concentration.

Ex.16 Why alloys do not oxidise (burn) readily at high temperature?

Sol. Because with the change in temperature their resistivity changes less rapidly.

Ex.17 Find the specific resistance of a wire of length 1.1m, 0.4 mm in diameter and having a total resistance of 4.2Ω.

Sol. $\therefore R = \frac{\rho \ell}{A} \Rightarrow \rho = \frac{RA}{\ell} \qquad \frac{RA}{\ell} = \frac{4.2 \times \pi d^2}{1.1 \times 4}$

Here $A = \pi r^2$ and $r = \frac{d}{2}$, $d = 0.4 \times 10^{-3} \text{m}$

$\therefore \rho = 48 \times 10^{-8} \Omega \text{ metre.}$

Ex.18 A wire of resistance 4 Ω is redrawn by pulling it doubled, what is its new resistance?

Sol. Let L is the length of the wire before stretching, L' is the length of the wire after stretching, A is the area of the wire before stretching, A' is the area of the wire after stretching, R is the resistance of the wire before stretching,, R' is the resistance of the wire after stretching.

Since the volume of the wire remains constant,

$\therefore AL = A'L'$

$\Rightarrow L' = 2L$

So, $A' = \frac{AL}{L} = \frac{AL}{2L} = \frac{A}{2}$

$\Rightarrow \frac{R'}{R} = \frac{\rho L}{A} \times \frac{A}{1L} = \frac{A}{A} \times \frac{L'}{L} = \frac{A}{A/2} \times \frac{2L}{L}$

So, $\frac{R'}{R} = 4$

$R' = 4R$


$R = 4\Omega$

Hence $R' = 4 \times 4 = 16 \Omega$.

Combination of Resistances

◆ Series Combination

When two or more resistances are joined end-to-end so that the same current flows through each of them, they are said to be connected in series.



When a series combination of resistances is connected to a battery, the same current (I) flows through each of them.

- **Law of combination of resistances in series:** The law of combination of resistances in series states that when a number of resistances are connected in series, their equivalent resistance is equal to the sum of the individual resistances. Thus, if R_1 , R_2 , R_3 ..., etc. are combined in series, then the equivalent resistance (R) is given by,

$$R = R_1 + R_2 + R_3 + \dots \quad \dots(i)$$

- **Derivation of mathematical expression of resistances in series combination:** Let R_1 , R_2 and R_3 be the resistances connected in series, I be the current flowing through the circuit, i.e., passing through each resistance, and V_1 , V_2 and V_3 be the potential difference across R_1 , R_2 and R_3 , respectively. Then, from Ohm's law,

$$V_1 = IR_1, V_2 = IR_2 \text{ and } V_3 = IR_3 \quad \dots(ii)$$

If V is the potential difference across the combination of resistances then,

$$V = V_1 + V_2 + V_3 \quad \dots(iii)$$

If R is the equivalent resistance of the circuit, then

$$V = IR \quad \dots(iv)$$

Using Eqs. (i) to (iv) we can write,

$$IR = V = V_1 + V_2 + V_3$$

$$= IR_1 + IR_2 + IR_3$$

$$\text{or, } IR = I(R_1 + R_2 + R_3)$$

$$\text{or, } R = R_1 + R_2 + R_3$$

Therefore, when resistances are combined in series, the equivalent resistance is higher than each individual resistance.


- **Some results about series combination:**

- (i) When two or more resistors are connected in series, the total resistance of the combination is equal to the sum of all the individual resistances.
- (ii) When two or more resistors are connected in series, the same current flows through each resistor.
- (iii) When a number of resistors are connected in series, the voltage across the combination (i.e. voltage of the battery in the circuit), is equal to the sum of the voltage drop (or potential difference) across each individual resistor.


$$(iv) R_{eq} > R_{max}$$

◆ Parallel Combination

When two or more resistances are connected between two common points so that the same potential difference is applied across each of them, they are said to be connected in parallel.



Parallel Circuit



Equivalent Circuit of Parallel Combination

When such a combination of resistance is connected to a battery, all the resistances have the same potential difference across their ends.

• Derivation of mathematical expression of parallel combination:

Let V be the potential difference across the two common points A and B. Then, from Ohm's law

$$\text{Current passing through } R_1, I_1 = V/R_1 \quad \dots(i)$$

$$\text{Current passing through } R_2, I_2 = V/R_2 \quad \dots(ii)$$

$$\text{Current passing through } R_3, I_3 = V/R_3 \quad \dots(iii)$$

If R is the equivalent resistance, then from Ohm's law, the total current flowing through the circuit is given by,

$$I = V/R \quad \dots(iv)$$

$$\text{and } I = I_1 + I_2 + I_3 \quad \dots(v)$$

Substituting the values of I , I_1 , I_2 and I_3 in eq. (v),

$$\frac{V}{R} = \frac{V}{R_1} + \frac{V}{R_2} + \frac{V}{R_3} \quad \dots(vi)$$

Cancelling common V term, one gets

$$\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3}$$


The equivalent resistance of a parallel combination of resistance is less than each of all the individual resistances.

• Important results about parallel combination:

- (i) Total current through the circuit is equal to the sum of the currents flowing through it.
- (ii) In a parallel combination of resistors the voltage (or potential difference) across each resistor is the same and is equal to the applied voltage i.e. $V_1 = V_2 = V_3 = V$
- (iii) Current flowing through each resistor is inversely proportional to its resistances, thus higher the resistance of a resistor, lower will be the current flowing through it.
- (iv) $R_{eq} < R_{min}$


Ex.19 Draw a schematic diagram of a circuit consisting of a battery of three cells of 2 V each, a 5Ω , 8Ω and 12Ω resistors, and a plug key, all connected in series.

Sol. Three cells of potential 2 V, each one of these cells are connected in series therefore the potential difference of the combined battery will be $2\text{ V} + 2\text{ V} + 2\text{ V} = 6\text{ V}$. The following circuit diagram shows three resistors of resistances 5Ω , 8Ω and 12Ω respectively connected in series and a battery of potential 6 V and a plug key which is closed means the current is flowing in the circuit.



Ex.20 Redraw the circuit of above question, putting in an ammeter to measure the current through the resistors and a voltmeter to measure the potential difference across the 12Ω resistor. What would be the readings in the ammeter and the voltmeter?

Sol.



According to Ohm's law,

$$V = IR,$$

Now Potential difference (V) = 6 V

Let current flowing through the circuit = I

Resistance of the circuit, $R = 5 + 8 + 12 = 25\Omega$

Since $I = V/R$

$$= 6 / 25$$

$$= 0.24\text{ A}$$

Potential difference across 12Ω resistor = V_1

Current flowing through the 12Ω resistor, $I = 0.24\text{ A}$

Therefore, using Ohm's law, we obtain

$$V_1 = IR$$

$$= 0.24 \times 12 = 2.88\text{ V}$$

Therefore, the reading of the ammeter will be 0.24 A.

And the reading of the voltmeter will be 2.88 V.

Ex.21 Judge the equivalent resistance when the following are connected in parallel –

(a) 1Ω and $10^6\Omega$

(b) 1Ω , $10^3\Omega$ and $10^6\Omega$.

Sol. (a) Since 1Ω and $10^6\Omega$ are connected in parallel then the equivalent resistance R will be

$$\frac{1}{R} = \frac{1}{1} + \frac{1}{10^6}$$

$$\Rightarrow R = \frac{10^6}{10^6 + 1} = \frac{10^6}{10^6} = 1 \Omega$$

$\therefore$ Equivalent resistance is 1Ω

(b) 1Ω and $10^3 \Omega$ and $10^6 \Omega$ are in parallel

$$\frac{1}{R} = \frac{1}{1} + \frac{1}{10^3} + \frac{1}{10^6}$$

$$\frac{1}{R} = \frac{10^6 + 10^3 + 1}{10^6}$$

$$\frac{1}{R} = \frac{1001001}{1000000} \Rightarrow R = 0.999 \Omega$$

Therefore, equivalent resistance = 0.999Ω

Ex.22 An electric lamp of 100Ω , a toaster of resistance 50Ω , and a water filter of resistance 500Ω are connected in parallel to a 220 V source. What is the resistance of an electric iron connected to the same source that takes as much current as all three appliances, and what is the current through it?

Sol. Resistance of electric lamp (R_1) = 100Ω

Resistance of toaster (R_2) = 50Ω

Resistance of water filter (R_3) = 500Ω

Potential difference of the source, $V = 220 \text{ V}$

Since all the resistances are in parallel then the equivalent resistance R of the circuit will be

$$\begin{aligned} \frac{1}{R} &= \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} = \frac{1}{100} + \frac{1}{50} + \frac{1}{500} \\ &= \frac{5 + 10 + 1}{500} = \frac{16}{500} \end{aligned}$$

$$R = \frac{500}{16}$$

According to Ohm's law

$$I = \frac{V}{R}$$

Where current flowing through the circuit = I

$$I = \frac{220}{\frac{500}{16}} = \frac{220 \times 16}{500} = 7.04 \text{ A}$$

All the three given appliances are drawing 7.04 A of current.

Therefore, current drawn by an electric iron connected to the same source of potential $220 \text{ V} = 7.04 \text{ A}$

Let R' be the resistance of the electric iron. According to Ohm's law,

$$V = IR$$

$$R' = \frac{V}{I} = \frac{220}{7.04} = 31.25 \Omega$$

Therefore, the resistance of the electric iron is 31.25Ω and the current flowing through it is 7.04 A .

Heating Effect of Electric Current

When the ends of a conductor are connected to a battery, then free electrons move with drift velocity and electric current flows through the wire. These electrons collide continuously with the positive ions of the wire and thus the energy taken from the battery is dissipated. To maintain the electric current in the wire, energy is taken continuously from the battery. This energy is transferred to the ions of the wire by the electrons. This increases the thermal motion of the ions, as a result the temperature of the wire rises. The effect of electric current due to which heat is produced in a wire when current is passed through it is called heating effect of current or Joule's heating. In 1841 Joule found that when current is passed through a conductor the heat produced across it is:

- (i) Directly proportional to the square of the current through the conductor i.e. $H \propto I^2$
- (ii) Directly proportional to the resistance of the conductor i.e. $H \propto R$
- (iii) Directly proportional to the time for which the current is passed i.e. $H \propto t$

Combining the above three equations we have,

$$H \propto I^2 R t$$

$$\text{or } H = \frac{I^2 R t}{J} \text{ (in calorie)}$$

Where J is called Joule's mechanical equivalent of heat and has a value of $J = 4.18 \text{ J cal}^{-1}$. The above equation is called Joule's law of heating.

In some cases, heating is desirable, while in many cases, such as electric motors, generators or transformers, it is highly undesirable. Some of the devices in which heating effect of an electric current is desirable, are incandescent lamps, toasters, electric irons and stoves. The tungsten filament of an incandescent lamp operates at a temperature of 2700°C . Here, we see electrical energy being converted into both heat and light energy.

◆ Electric Energy

The fact that conductors offer resistance to the flow of current, means that work must be continuously done to maintain the current. The role of resistance in electrical circuits is analogous to that of friction in mechanics. The amount of work done by current I , flowing through a wire of resistance R during the time t is calculated by -

$$W = QV$$

but as $Q = I \times t$

Therefore, the amount of work done, W is

$$W = V \times I \times t$$

By substituting the expression for V from Ohm's law,

$$V = IR$$

we finally get $W = I^2 R t$

This shows that the electrical energy dissipated or consumed depends on the product of the square of the current I , flowing through the resistance R and the time t .

Ex.30 Why does the cord of an electric heater not glow while the heating element does, when connected to the mains?

Sol. The wires of the connecting cord of electric heater are made of Cu, negligible heat is produced in it, due to its extremely low resistance. The heating element of an electric heater is made up of nichrome wire. It glows because large amount of heat is produced due to its high resistance.

Ex.31 Compute the heat generated while transferring 96000 coulomb of charge in one hour through a potential difference of 50 V.

Sol. The amount of heat (H) produced is given by the Joule's law of heating as $H = VIt$
where, Voltage, $V = 50$ V

Time, $t = 1$ h = $1 \times 60 \times 60$ s = 3600 s

Amount of current, $I = q/t = 96000 / 3600 = (80/3)$ A

$$H = 50 \times \frac{80}{3} \times 60 \times 60 = 4.8 \times 10^6 \text{ J}$$

Therefore, the heat generated is 4.8×10^6 J.

Ex.32 An electric iron of resistance 20Ω takes a current of 5 A. Calculate the heat developed in 30 s.

Sol. The amount of heat (H) produced is given by the Joule's law of heating as $H = VIt$
where, Current, $I = 5$ A

Time, $t = 30$ s

Voltage, $V = \text{Current} \times \text{Resistance} = 5 \times 20 = 100$ V

$$H = 100 \times 5 \times 30 = 1.5 \times 10^4 \text{ J}$$

Therefore, the amount of heat developed in the electric iron is 1.5×10^4 J.

- Commercial unit of electrical energy (Kilowatt - hour): The S.I. unit of electrical energy is joule and we know that for commercial purposes we use a bigger unit of electrical energy which is called "kilowatt - hour". One kilowatt - hour is the amount of electrical energy consumed when an electrical appliance having a power rating of 1 kilowatt and is used for 1 hour.

- Relation between kilowatt hour and Joule: Kilowatt-hour is the energy supplied by a rate of working of 1000 watts for 1 hour.

$$1 \text{ kilowatt-hour} = 3600000 \text{ Joules}$$

$$1 \text{ kWh} = 3.6 \times 10^6 \text{ J}$$

◆ Electric Power

The rate at which electric energy is dissipated or consumed, is termed as electric power. The power P is given by,

$$P = W/t = I^2 R$$

The unit of electric power is watt, which is the power consumed when 1A of current flows at a potential difference of 1 V.

- **Unit of power:** The S.I. unit of electric power is 'watt' which is denoted by the letter W. The power of 1 watt is a rate of working of 1 joule per second.

A bigger unit of electric power is kilowatt.

$$1 \text{ kilowatt (kW)} = 1000 \text{ watt.}$$

Power of an agent is also expressed in horse power (hp).

$$1 \text{ hp} = 746 \text{ watt}$$

- **Formula for calculating electric power:**

$$\text{We know, Power, } P = \frac{\text{Work}}{\text{Time}}$$

and Work, $W = V \times I \times t$

$$P = \frac{V \times I \times t}{t} \Rightarrow P = V \times I$$

Power P in terms of I and R :

Now from Ohm's law we have, $\frac{V}{I} = R$

$$V = I \times R$$

$$P = I \times R \times I$$

$$P = I^2 \times R$$

Power P in terms of V and R:

We know, $P = V \times I$

From Ohm's law $I = \frac{V}{R}$

$$P = V \times \frac{V}{R} \Rightarrow P = \frac{V^2}{R}$$

Ex.33 What determines the rate at which energy is delivered by a current?

Sol. The rate of consumption of electric energy in an electric appliance is called electric power. Hence, the rate at which energy is delivered by a current is the power of the appliance.

Ex.34 An electric motor takes 5 A from a 220 V line. Determine the power of the motor and the energy consumed in 2 h.

Sol. Power of motor = $VI = 220 \times 5 = 1100 \text{ W}$

Energy consumed in 2 h = Power $\times$ time = $1100 \text{ W} \times 2 \times 60\text{s} \times 60\text{s} = 1100 \text{ W} \times 7200\text{s} = 7.92 \times 10^6 \text{ J}$

- Calculation of Electric bill:

Energy consumed by electric appliances is given by the formula.

$$\text{Electric energy (in kWh)} = \frac{\text{no. of appliances} \times \text{Power of appliances (in watt)} \times \text{Time (in hour)}}{1000}$$

◆ Power-Voltage Rating of Electrical Appliances

Every electrical appliance like an electric bulb, radio or fan has a label or engraved plate on it which tells us the voltage (to be applied) and the electrical power consumed by it. For example, if we look at a particular bulb in our home, it may have the figures 220 V, 100 W written on it. Now, 220 V means that this bulb is to be used on a voltage of 220 volts and 100 W which means, it has a power consumption of 100 watts or 100 joules per second, when supplied a voltage of 220 volt.

◆ Applications of Heating Effect of Current

Domestic electrical appliances such as electric bulb, electric iron, geyser, room heater etc work on heating effect of current and are rated in terms of voltage and wattage. The coils of these devices are made of a material of a very high resistance, (for instance, nichrome or tungsten) such that when a current passes through the coil, heat is generated. Generally the potential difference applied to the electrical appliance is the same as that of the mains i.e. 220 - 230 V in India and 110 V in U.S.A., Canada etc.

Note: The most popular heating element nichrome is an alloy of 60% nickel, 25% iron and 15% chromium. It has low resistance and high melting point.

◆ Electric Fuse

An electric fuse is an easily fusible wire of short length put into an electrical circuit for protection purposes. It is arranged to melt ("blow") at a definite current.

It is an alloy of lead and tin (37% lead + 63% tin). It has a high resistivity and low melting point. As soon as the safe limit of current exceeds, the fuse "blows" and the electric circuit is cut off.

EXERCISE-1

➤ Very Short Answer Type Questions

- Q.1** What is represented by joule/coulomb?
- Q.2** What is the function of electric switch in electric circuit?
- Q.3** A wire of resistivity 10 ohm meter is stretched to double its length. What is its new resistivity?
- Q.4** What is the SI unit of Power?
- Q.5** A charge of 2C moves between two plates, maintained at a p.d of 1V. What is the energy acquired by the charge?

➤ Short Answer Type Questions – Type I

- Q.6** Draw a schematic diagram of a typical electric circuit comprising a cell, an electric bulb, an ammeter and a plug key.
- Q.7** What makes the electric charge to flow in a conducting metallic wire?
- Q.8** Two wires of same material and same length have radii r_1 and r_2 , compare their resistances.
- Q.9** A wire of length L and resistance R is stretched so that its length is doubled. How will its
(a) Resistance change
(b) Resistivity change?
- Q.10** Calculate the electrical energy consumed by a 1200W toaster in 20 minutes.

➤ Short Answer Type Questions – Type II


- Q.11** Write any three properties of charges.
- Q.12** If '2q' coulombs of charge travel through a conductor of length ' ℓ ' m with a velocity of ' v ' m/s, what is the current flowing through the conductor?

- Q.13** A TV set shoots out a beam of electrons. The beam current is $10\mu\text{A}$.
(a) How many electrons strike the TV screen in each second?
(b) How much charge strikes the screen in a minute?
- Q.14** (a) Why do copper or aluminium wires generally used for electrical transmission and distribution purposes?
(b) Two wires, one of copper and other of manganin, have equal lengths and equal resistances. Which wire is thicker? Given that resistivity of copper is lower than that of manganin.

- Q.15** Two conductors X and Y of circular cross-section have radii in the ratio of 1: 2, length in the ratio 1: 3 and resistivities in the ratio of 3: 1, then in which ratio their resistances are?

- Q.16** Given n resistors each of resistance r . How will you combine them to get the
(i) Maximum and
(ii) Minimum effective resistance? What is the ratio of the maximum to minimum resistance?

- Q.17** Three resistances are connected as shown in diagram. Through the resistance 5 ohms, a current of 1 ampere is flowing:



- (i) What is the current through the other two resistors?
(ii) What is the potential difference (p.d.) across AB and across AC?
(iii) What is the total resistance?
- Q.18** Explain heating effect of current in brief and deduce the formula used..

Q.19 A household uses the following electrical appliances:

- (i) Refrigerator of rating 400 W for ten hours every day.
- (ii) Two electric fans of rating 80 W each for twelve hours every day.
- (iii) Six electric tubes of rating 18 W each for 6 hours every day.

Calculate the electricity bill of the household for the month of June if the cost per unit of electric energy is Rs. 3.00.

Q.20 Two lamps, one rated 60 W at 220 V and the other 40 W at 220 V, are connected in parallel to the electric supply at 220 V.

- (a) Draw a circuit diagram to show the connections.
- (b) Calculate the current drawn from the electric supply.
- (c) Calculate the total energy consumed by the two lamps together when they operate for one hour.

➤ Long Answer Type Questions

Q.21 Write any five differences between static and current electricity?

Q.22 What is potential difference? Define its SI unit. Which instrument is used to measure it and how it is used?

Q.23 Write activities that can be performed to show that in a series combination of resistors the current is same in every part of the circuit or same current passes through each resistor.



Q.24 Derive the equation for resultant resistance of resistors in series.

Q.25 Derive the equation for resultant resistance of resistors in parallel.


➤ Practical & Value Based Type Questions

Q.26 Two students perform the experiments on series and parallel combinations of two given resistors R_1 and R_2 and plot the following V-I graphs.

Which of the two diagrams correctly represent the labels 'series and parallel' on the plotted curves? Justify your answer.



Q.27 The figures given below show the readings of a milliammeter and a voltmeter connected in an electrical circuit. Assuming that the instruments do not have any zero error, what will be the correct readings of the milliammeter and voltmeter?




Q.28 In the evening Seema parked her car inside her garage and left for vacation of 15 days. On returning, she tried to start the car but the car did not start. She checked the fuel and engine oil but that were also full. So, she called a mechanic. After examining the car and mechanic connected the charged battery with that of car battery. Then the car started at once. On the basis of above passage, answer the following questions

- (a) In which combination, the batteries are connected?
- (b) Draw a circuit diagram for the above said combination.
- (c) Identify the moral values we get from the passage.

EXERCISE-2

- Q.1** When a body is negatively charged by friction, it means
 (A) the body has acquired excess of electrons
 (B) the body has acquired excess of protons
 (C) the body has lost some electrons
 (D) the body has lost some neutrons
- Q.2** When the distance between the charged particles is halved, the force between them becomes
 (A) One-fourth (B) Half
 (C) Double (D) Four times
- Q.3** How many electrons in 1 s constitute a current of 1 A?
 (A) 6.25×10^{18} (B) 6.25×10^{12}
 (C) 6.25×10^{11} (D) 6.25
- Q.4** If a charged body attracts another body, the charge on the other body
 (A) must be negative
 (B) must be positive
 (C) must be zero
 (D) may be negative or positive or zero
- Q.5** A suitable unit for expressing the strength of electric field is
 (A) V/C (B) C/m (C) N/C (D) C/N
- Q.6** A particle of mass m and charge q is placed at rest in a uniform electric field E and then released. The kinetic energy attained by the particle after moving a distance y is
 (A) qEy^2 (B) qE^2y
 (C) qEy (D) q^2Ey
- Q.7** The number of electrons to be put on a spherical conductor of radius $0.1m$ to produce an electric field of $0.036N/C$ just above its surface is
 (A) 2.7×10^5 (B) 2.6×10^5
 (C) 2.5×10^5 (D) 2.4×10^5
- Q.8** Deuteron and α -particle are put $1\text{\AA}$ apart in air. Magnitude of intensity of electric field due to deuteron at α -particle is
 (A) Zero
 (B) 2.88×10^{11} newton/coulomb
 (C) 1.44×10^{11} newton/coulomb
 (D) 1.44×10^{11} newton/coulomb
- Q.9** What is not true for electric charge :
 (A) Electric charge is scalar quantity
 (B) Charge on a body may be +ve or -ve
 (C) S.I. unit of charge is coulomb
 (D) One coulomb is charge of one electron
- Q.10** A metallic sphere has a charge of $10\mu C$. A unit negative charge is brought from A to B both 100 cm away from the sphere but A being east of it while B being on west. The net work done is
 (A) Zero (B) 2/10 joule
 (C) -2/10 joule (D) -1/10 joule
- Q.11** 1 Coulomb is equal to
 (A) 1 amp $\times$ 1 sec (B) 1 amp / 1 sec
 (C) 1 joule $\times$ 1 amp (D) 1 joule / 1 sec
- Q.12** One ampere equals
 (A) $10^6 \mu A$ (B) $10^{-6} \mu A$
 (C) $10^{-3} \mu A$ (D) $10 \mu A$
- Q.13** If I is the current through a wire and 'e' is the charge of electron then the number of electrons in t seconds will be given by
 (A) $\frac{Ie}{t}$ (B) e/It
 (C) It/e (D) Ite
- Q.14** Figure shows current in a part of electrical circuit, then the value of current is-




- (A) 1.7 A (B) 3.7 A
 (C) 13 A (D) 1.0 A

- Q.15** The unit of resistivity is
 (A) ohm (B) ohm mete
 (C) ohm meter⁻¹ (D) mho metre⁻¹

- Q.16** If a wire of resistance $1\ \Omega$ is stretched to double its length, then the resistance will become
 (A) $\frac{1}{2}\ \Omega$ (B) $2\ \Omega$ (C) $\frac{1}{4}\ \Omega$ (D) $4\ \Omega$

- Q.17** In the given circuit, the effective resistance between points A and C will be




- (A) $\frac{3}{2}R$ (B) $6R$ (C) $\frac{2}{3}R$ (D) $3R$

- Q.18** A wire of resistance R is cut into n equal parts. These parts are then connected in parallel. The equivalent resistance of combination will be
 (A) nR (B) R/n (C) n/R (D) R/n^2

- Q.19** A piece of wire of resistance $4\ \Omega$ is bent through 180° at its mid point and the two halves are twisted together, then resistance is
 (A) $1\ \Omega$ (B) $2\ \Omega$ (C) $5\ \Omega$ (D) $8\ \Omega$

- Q.20.** Calculate the equivalent resistance between P & Q of the network. Show in the figure given.



- (A) $8\ \Omega$ (B) $4\ \Omega$ (C) $6\ \Omega$ (D) $9\ \Omega$

- Q.21** Which of the following effects of current does not depend on direction of current?
 (A) Lighting and chemical effect
 (B) Heating and lighting effect
 (C) Heating and magnetic effect
 (D) Magnetic and chemical effect

- Q.22** Two heater wires of equal length are first connected in series and then in parallel with a battery. The ratio of heat produced in the two cases is
 (A) 2: 1 (B) 1: 2 (C) 4: 1 (D)* 1: 4

- Q.23** The resistance of a conductor is reduced to half its initial value. In doing so the heating effects in the conductor will become
 (A) Half (B) One-fourth
 (C) Four times (D) Double

- Q.24** Laws of heating are given by
 (A) Faraday (B) Joule
 (C) Ohm (D) Maxwell

- Q.25** Which of the following statement is false?
 (A) Heat produced in a conductor is proportional to its resistance
 (B) Heat produced in a conductor is proportional to the square of the current
 (C) Heat produced in a conductor is proportional to charge
 (D) Heat produced in a conductor is proportional to the time for which current is passed


- Q.26** The filament of an electric bulb is made of tungsten because
 (A) Its resistance is negligible
 (B) It is cheaper
 (C) Its melting point is high
 (D) Its filament is easily made

- Q.27** Electric power is transmitted over long distance through conducting wires at very high voltage because
 (A) The power losses are reduced to minimum
 (B) Signal of high voltage travel faster
 (C) High voltage can be stepped down to desired levels easily
 (D) None of these

EXERCISE-3


- Q.1** In the given circuit diagram, the value of resistance X in ohm when the bridge is balanced will be

[Raj./ NTSE Stage-I/2005]



- (A) 4 (B) 8
(C) 10 (D) 12


- Q.2** In the following figure, the equivalent resistance between the points A and B in ohm will be [Raj./ NTSE Stage-I/2005]



- (A) 1 (B) 2
(C) (D) 9

- Q.3** In the circuit shown in figure, the current flowing through 5Ω resistance is

[Raj./ NTSE Stage-I/2006]



- (A) 0.5A (B) 0.6A
(C) 0.9 A (D) 1.5 A

- Q.4** The material of heating element in an electric press has

[Raj./NTSE Stage-I/2007]

- (A) high melting point and high specific resistance
(B) high melting point and low specific resistance
(C) low melting point and high specific resistance
(D) low melting point and high specific resistance

- Q.5** A uniform wire when connected directly across a 220V line produces heat H per second. If the wire is divided into n -parts and all parts are connected in parallel across a 220 V line, the heat produced per second will be

[Haryana/NTSE Stage-I/2014]

- (A) H/n (B) H/n^2
(C) n^2H (D) nH

- Q.6** One Volt is equal to

[Raj./ NTSE Stage-I/2013]

- (A) 1 Joule
(B) 1 Newton/Coulomb
(C) 1 Joule/Coulomb
(D) 1 Coulomb/Newton

- Q.7** On a bulb it is written as 220 Volt and 60 watt. Find out the resistance of the bulb and the value of the current flowing through it

[Raj./ NTSE Stage-I/2013]

- (A) 806.66 ohm / 0.27 ampere
(B) 500 ohm / 2 ampere
(C) 200 ohm / 4 ampere
(D) 100 ohm / 1 ampere


- Q.8** A comb run through one's dry hair attracts small bits of paper. This is due to

[Delhi/ NTSE Stage-I/2014]

- (A) comb is a good conductor
(B) paper is a good conductor
(C) The atoms in the paper gets polarised by the charged comb
(D) the comb possesses magnetic properties

- Q.9** The value of current I and voltage V in the given circuit will be

[Raj./ NTSE Stage-I/2014]



- (A) 2A, 4V (B) 4A, 2V
(C) 1A, 2V (D) 2A, 1V


- Q.10** When 1 J of work is done to move a charge of 1 C from one point to another point then the potential difference between two points in a given circuit will be

[Raj./NTSE Stage-I/2015]

- (A) 1 V (B) 4V
(C) 8 V (D) zero.

- Q.11** The value of equivalent resistance between the point A and B in the given circuit will be


[Raj./NTSE Stage-I/2015]



- (A) 6 R (B) $\frac{4R}{11}$
(C) $\frac{11R}{4}$ (D) $\frac{R}{6}$

- Q.12** Find the reading of the ammeter in the circuit given below

[Raj./ NTSE Stage-II/2015]



- (A) $\frac{V}{2R}$ (B) $\frac{3V}{4R}$
(C) $\frac{2V}{7R}$ (D) $\frac{11V}{R}$

- Q.13** A certain household has consumed 200 units of energy during a month. Its value in joules will be [Raj./NTSE Stage-I/2015]

- (A) 3.6×10^{10} (B) 7.2×10^{10}
(C) 3.6×10^8 (D) 7.2×10^8

- Q.14** In a neon gas discharge tube, every second 4.8×10^{18} Ne^+ ions move towards the right through a cross-section of the tube, while 'n' electrons move to the left in the same time. If the current in the tube is 1.12 amperes towards the right, n is equal to (given $e = 1.6 \times 10^{-19}$ coulomb)

[Raj./ NTSE Stage-II/2016]

- (A) 1.8×10^{18} (B) 2.2×10^{18}
(C) 2.4×10^{19} (D) 2.8×10^{19}

- Q.15** Four 20Ω resistors are connected in series and the combination is connected to a 20 V emf device. The potential difference across any one of the resistors is -

[Raj./NTSE Stage-I/2016]

- (A) 5V (B) 2V
(C) 4V (D) 20V

- Q.16** Of the following, the copper conductor that has the least resistance is

[Raj./ NTSE Stage-I/2016]

- (A) thin, long and hot
(B) thick, short and cool
(C) thick long and hot
(D) thin, short and cool

- Q.17** Three 6Ω resistors are connected in parallel and the combination is connected to a 15 V battery. The current through any one of the resistors will be

[Raj./NTSE Stage-I/2017]


- (A) 2.5 A (B) 2.0 A
(C) 5 A (D) 10 A

- Q.18** The electrical resistivity of a conducting wire is K. If its length and area of cross section are doubled then the new resistivity of the wire will be

[Raj./NTSE Stage-I/2018]

- (A) K (B) 2K
(C) K/2 (D) K/4


- Q.19** What is the equivalent resistance of the given circuit between points A and B?



[Raj./NTSE Stage-I/2018]

- (A) $10\ \Omega$ (B) $4\ \Omega$
 (C) $\frac{14}{3}\ \Omega$ (D) $\frac{17}{6}\ \Omega$
- Q.20** 4 bulbs rated 100 W each, operate for 6 hours per day. What is the cost of the energy consumed in 30 days at the rate of Rs. 5 /kWh? [Raj./NTSE Stage-I/2018]
- (A) Rs. 360 (B) Rs. 90
 (C) Rs. 120 (D) Rs. 400
- Q.21** A constant current I flows in a horizontal wire in the plane of the paper from West to East as shown in the figure. The direction of magnetic field at a point will be South to North

[DEL./NTSE Stage-I/2019]



- (A) directly above the wire
 (B) directly below the wire
 (C) at a point located in the plane of the paper, on the north side of the wire.
 (D) at a point located in the plane of the paper, on the south side of the wire.

- Q.22** If the current through a resistor is increased by 50%, the increase in power dissipated will be (assume the temperature remains constant) [DELHI/NTSE Stage-I/2019]
- (A) 225% (B) 200%
 (C) 250% (D) 125%

- Q.23** Conductivity of superconductors is: [MP./NTSE Stage-I/2019]
- (A) Infinite (B) Very large
 (C) Very Small (D) Zero

- Q.24** 1k Wh equal to: [MP./NTSE Stage-I/2019]
- (A) $3.6 \times 10^4\ \text{J}$ (B) $3.6 \times 10^5\ \text{J}$
 (C) $3.6 \times 10^6\ \text{J}$ (D) $3.6 \times 10^7\ \text{J}$

- Q.24** In a Helium gas discharge tube every second $40 \times 10^{18}\ \text{He}^+$ (ions) move towards the right through a cross – section of the tube, while n electrons move to the left in the same time. If the current in the tube is 8A to towards right then $n = ?$ [Maharashtra/ NTSE Stage-I/2019]

- (A) 3×10^{18} (B) 3×10^{19}
 (C) 3×10^{20} (D) 3×10^{21}

ANSWER KEY

EXERCISE - 1

3. 40ω
5. 2 J
8. $\frac{R_1}{R_2} = \left(\frac{r_2}{r_1}\right)^2$
10. $1.44 \times 10^6 \text{ J}$
12. $2qv/\ell$
13. (a) 6.25×10^{13} (b) $6 \times 10^{-4} \text{ C}$
15. 4:1
16. (i) series (ii) parallel (iii) n^2
17. (i) $I_1 = 0.6 \text{ A}$, $I_2 = 0.4 \text{ A}$. (ii) P.D. across AB = 5 V. P.D. across AC = 11 V
(iii) Total circuit resistance = 11 Ω .
19. Rs 591 (Approx.)
20. 0.45 A, $3.6 \times 10^5 \text{ J}$
26. Both are correct because $\Delta V/\Delta I = \text{resistance}(R)$ and $\Delta I/\Delta V = 1/R$
Series combination gives high resistance and parallel combination gives low resistance
27. Correct reading of ammeter = 130 Ma
Correct reading of voltmeter = 1.5 V

EXERCISE - 2

Ques.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Ans.	A	D	A	D	C	C	C	C	D	A	A	A	C	A	B
Ques.	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
Ans.	D	C	D	A	A	B	D	D	B	C	C	A	C	C	A

EXERCISE - 3

Ques.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Ans.	B	A	B	B	B	C	A	C	A	A	C	B	D	B	A
Ques.	16	17	18	19	20	21	22	23	24						
Ans.	B	A	A	B	A	B	D	A	C						

CHEMICAL REACTIONS AND EQUATIONS

Chapter Outline


- ✧ Physical Change
- ✧ Chemical Change
- ✧ Chemical Reactions
- ✧ Chemical Equations
- ✧ Balancing of Equations
- ✧ Types of Chemical Reactions
 - ◆ Combination Reaction
 - ◆ Decomposition Reaction
 - ◆ Displacement Reaction
 - ◆ Double Displacement Reaction
 - ◆ Redox Reaction
- ✧ Oxidation Number
- ✧ Oxidation & Reduction in Day to Day Life



Experimentation




Chemical reactions



Chemical reactions

MIND MAP

CHEMICAL REACTIONS & EQUATIONS



CHEMICAL REACTIONS AND EQUATIONS

Introduction

Chemistry is defined as the branch of science which deals with the study of matter, dealing with the composition, structure, properties and changes occurring in matter.

Changes occurring in matter may be physical changes like melting of wax, freezing of water or may be chemical changes. In this chapter we will discuss chemical changes occurring in matter.

Physical Change

A change which alters some specific physical properties of matter, like its state, texture, magnetic or electrical condition or its colour, without causing any change in the composition of the substance is called physical change.

e.g.: Boiling of water, dissolving of salt in water, ringing of electric bell etc.

Chemical Change

A change which alters the specific properties of a material by bringing a change in its molecular composition is called a chemical change.

e.g.: Formation of curd, burning of wood, etc.

Chemical Reactions

The process involving a chemical change is called a chemical reaction.

◆ Process Involved in Chemical Reactions

During chemical reaction, a rearrangement of atoms takes place between the reacting substances to form new substances having entirely different properties.

Chemical reactions involve breaking of chemical bonds present in the reactant molecules and forming new chemical bonds to give the products. Thus, a chemical reaction simply involves rearrangement of atoms.

The substances which take part in a chemical reaction are called *reactants* while the new substances produced as a result of chemical reaction are called *products*.

Few examples of chemical reactions in daily life:

- (i) Burning piece of paper.
- (ii) Curdling of milk.
- (iii) Rusting of iron articles such as pan or nails.
- (iv) Fermentation of grapes.
- (v) Cooking of food.
- (vi) Digestion of food takes place in our body.
- (vii) We respire.
- (viii) Burning of candle wax.
- (ix) Burning of fuels like kerosene oil, petrol, LPG, etc.


◆ Characteristics of Chemical Reactions

There are some characteristics by which we can identify the occurrence of a chemical reaction. These are described below:


1. **Evolution of gas:** Some of the chemical reactions occur with the evolution of a gas or a mixture of gases.

For example:

- (i) Metals like zinc, magnesium, iron, etc., react with dilute hydrochloric acid with the evolution of hydrogen gas.




- (ii) When lead nitrate is heated, it gets decomposed to give a solid residue to lead monoxide with the evolution of nitrogen dioxide and oxygen gases.



- (iii) When calcium carbonate is treated with hydrochloric acid, gives calcium chloride with evolution of colourless and odourless gas carbon dioxide.




- (iv) When carbon disulphide is treated with oxygen, it form Carbon dioxide and Sulphur dioxide.




2. **Formation of precipitate:** Sometimes, when two solutions are mixed together, a solid gets separated from the solution. The solid, thus, separated is called the precipitate.

For example:

- (i) When an aqueous solution of sodium sulphate is mixed with a solution of barium chloride, a white precipitate of barium sulphate is produced.



- (ii) When the aqueous solutions of lead acetate and potassium iodide are mixed, a yellow precipitate of lead iodide is produced.





3. **Change in temperature:** Change in temperature is a necessary characteristic of a chemical reaction. On the basis of change in temperature, reactions are of two types –

- ♦ **Exothermic:** Exothermic reactions are those which are accompanied with release of energy.

For example:

- (i) When quicklime (calcium oxide) is treated with water in a beaker, a large quantity of heat is produced. As a result, the beaker becomes very hot. Such reactions in which heat is produced are called exothermic reactions.



- (ii) $\text{CH}_4\text{(g)} + 2\text{O}_2\text{(g)} \rightarrow \text{CO}_2\text{(g)} + 2\text{H}_2\text{(g)} + \text{Energy}$
Methane


- (iii) $\text{C}_6\text{H}_{12}\text{O}_6\text{(aq)} + 6\text{O}_2\text{(g)} \rightarrow 6\text{CO}_2\text{(g)} + 6\text{H}_2\text{O} + \text{Energy}$
Glucose

- (iv) $2\text{Mg(s)} + \text{O}_2\text{(g)} \rightarrow 2\text{MgO(s)} + \text{Heat}$
Magnesium oxide


- ♦ **Endothermic:** Endothermic reactions are those which are accompanied with absorption of energy.

For example:

- When ammonium chloride and barium hydroxide are mixed together in a test tube and the mixture is stirred with a glass rod, the bottom of the test tube becomes cold. In this reaction, heat is absorbed. Such a reaction is called an endothermic reaction.




- $\text{N}_2(\text{g}) + \text{O}_2(\text{g}) + \text{Heat} \rightarrow 2\text{NO}$
Nitric oxide




- Change in colour:** In some chemical reactions a change in colour is observed.

For example:

- When lead nitrate (colourless) is made to react with potassium iodide (colourless), yellow coloured ppt of lead iodide is formed.




- When iron (grayish black) is left exposed to air for a long time, the outer surface of iron gets covered with a brown coating of rust.




- Physical state:** In some reactions the physical state of the product becomes different from that of the reactant.

For example:

- Magnesium ribbon (crystalline) is burnt it changes to white powder of magnesium oxide.



- Coal burns to form carbon dioxide which is gaseous.




One or more of the above characteristics can certify that a chemical reaction is occurring.


♦ Conditions for a Chemical Reaction

The important conditions for a chemical change (chemical reaction) to take place is as follows:

- Close contact of reactants:** A chemical change (chemical reaction) occurs when the reactants molecules responsible for the change are brought closer to each other.
- Exchange of energy:** The chemical changes take place by exchange of energy in some form, i.e., evolution or absorption of energy in some form.
 - Heat:** A large number of chemical reactions show evolution or absorption of heat. For example, lime stone (CaCO_3) does not decompose at room temperature but on heating to about 1000°C , it undergoes a chemical change to form lime (CaO) and carbon dioxide (CO_2).




Similarly carbon burns in air to release heat



However in many of the exothermic reactions, heat is initially provided to reactant molecules in order to activate them for reaction.



- (ii) **Light:** Some chemical reactions take place with absorption or emission of light. The reactant molecules, on absorbing light, get activated and interact with one another to bring in a chemical change.

For example, a mixture of hydrogen (H_2) and chlorine (Cl_2) undergoes a chemical change to form hydrogen chloride (HCl) when exposed to sunlight.





Note: Similarly emission of light by fire flies is due to oxidation of luciferin protein of their wings. Phosphorus left out of burnt bodies after their cremation, reacts with atmosphere to glow in dark nights. The light so emitted is called cold light.

- (iii) **Electricity:** Several chemical reactions are noticed by passing electric current through the reactants either in the molten state or in the solution form or electricity is produced as a result of chemical change. For example, when electric current is passed in molten sodium chloride, two new substances-sodium and chlorine are obtained (electrolysis). In electrochemical cell say Daniel cell electricity is produced as a result of chemical reaction.



3. **Presence of catalyst:** Catalysts are the substances which influence the rate of a chemical reaction and remains unchanged in amount and chemical composition at the end of the reaction. Their physical state may change. There are several very slow chemical reactions which require certain catalysts to show an appreciable extent of reaction. For example, potassium chlorate decomposes appreciably to give oxygen and potassium chloride in the presence of manganese dioxide as catalyst. If manganese dioxide is not used, the decomposition occurs at a much higher temperature.



Ex.1 What is the role of a catalyst in a chemical reaction?

Sol. Catalyst helps in increasing the rate of reaction and hence reaction becomes fast.

Ex.2 On what basis is a chemical equation balanced?


Sol. A chemical equation is balanced on the basis of law of conservation of mass according to which mass can neither be created nor be destroyed.

Chemical Equations

The method of representing a chemical reaction with the help of symbols and formulae of the substances involved in it is known as a *chemical equation*.


There are two ways of representing a chemical reaction:

1. **In Terms of Words (Called Word Equation):** Zinc metal reacts with dilute sulphuric acid to form zinc sulphate and hydrogen gas. This reaction can be written in words as



This is known as the word equation.

2. In Terms of Symbols and Formulae (Called Symbol Equation): By putting the symbols and formulae of all the substances in the above word equation, we get the following chemical equation.



This is called a chemical equation.

◆ Writing of a Chemical Equation

To represent a chemical equation we write:

- The symbols and formulae of the reactants towards the left hand side (LHS) with plus sign between them.
- The symbols and formulae of the products towards the right hand side (RHS) with plus sign between them.
- An arrow ($\rightarrow$) sign is put between the reactants and the products, pointing from reactants towards products.


◆ Balanced and Unbalanced Chemical Equations

A balanced chemical equation has equal number of atoms of different elements in the reactants and products.

For example:

Zinc + Sulphuric acid $\rightarrow$ Zinc sulphate + Hydrogen

The above word equation may be represented by the following chemical equation



Let us examine the number of atoms of different elements on both sides of the arrow.

Element	Number of atoms in reactants (LHS)	Number of Atoms in products (RHS)
Zn	1	1
H	2	2
S	1	1
O	4	4


As the number of atoms of each element is the same on both sides of the arrow, the above equation is a balanced chemical equation.

An unbalanced or skeletal equation is one in which the number of atoms of different elements (one or more) is not equal on the two sides of the equation.

For example:

Potassium chlorate $\rightarrow$ Potassium chloride + Oxygen

The above word equation may be represented as



Let us examine the number of atoms of different elements in the unbalanced equation.

Element	Number of atoms in reactants (LHS)	Number of atoms in products (RHS)
K	1	1
Cl	1	1
O	3	2

As the number of atoms of K and Cl are equal on both sides but O-atoms are not equal on both sides, hence the above equation is unbalanced.

Balancing of a Chemical Equation

Balancing of a chemical equation means making the number of each element equal on both sides of the equation.

- ♦ **Importance of a Balanced Chemical Equation:** The importance of a balanced chemical equation lies in the fact that it satisfies the law of conservation of mass, i.e., in a chemical reaction, total mass of all the products is equal to the total mass of all the reactions.

- ♦ **Steps Involved in the Balancing of a Chemical Equation:**

The balancing of a chemical equation involves counting up the number of atoms of each element on both sides and trying to equalize them.

The following steps are involved in the balancing of a chemical equations:

Step-I: To write the word equations: Write the chemical equation in the form of a word equations. Keep the reactants on left hand side (LHS) and the products on the right hand side (RHS). Separated them by an arrow ($\rightarrow$) head of which points from the reactants towards the products.

Step-II: To write the skeletal chemical equation: Write down the symbols and formulae of the various reactants and products. This gives us the skeletal chemical equation.

Step-III: Enclose the formula of each reactant and product in a box. Do not change anything inside the boxes while balancing equation.

Step-IV: List the number of atoms of different elements on LHS (Reactants) and RHS (Products) in the unbalanced equation.

Step-V: Select the biggest formula to start balancing: It is often convenient to start balancing with the compound that contains the maximum number of atoms. It may be a reactant or a product.

Step-VI: To start balancing different elements: Having selected the compound with the biggest formula, first balance the element of this compound. Then balance other elements one by one. To balance the atoms of an element put a whole number coefficient before the formula of the compound.

If selection of the biggest formula appears inconvenient, balance the atoms of that element which occurs at minimum number of places on both sides of the equation. Atoms of the element which occur at maximum places are balanced last of all.

Step-VII: To check the correctness of the balanced equation: Finally, to check the correctness of the balanced equation, count atoms of each element on both sides of the equation. If the number of atoms of elements on both sides of the equation are equal then the equation is balanced.


The method of balancing chemical equation is called Hit and Trial Method as we make trial to balance the equation by using the smallest whole number coefficient.

Ex.3 Combustion of methane in oxygen to form carbon dioxide and water.

Sol. Step-I: Writing the equation in word form:

Methane + Oxygen $\rightarrow$ Carbon dioxide + Water


Step-II: Writing the skeletal equation:



Step-III: Selecting the element which occurs at minimum number of places, i.e. C or H.

No. of C-atoms is same on LHS & RHS.

No of H-atoms is 4 on LHS and 2 H-atoms of RHS. To balance H-atoms, multiply H_2O on RHS by 2, we get,



Step-IV: To balance O-atoms: There are 4 O-atoms on RHS and 2 O-atoms on LHS. To balance O-atoms, multiply O_2 on LHS by 2, we get,
 $CH_4 + 2O_2 \rightarrow CO_2 + 2H_2O$

Step-V: To check the corrections of the balanced equation:

Element	Number of atoms on LHS	Number of atoms on RHS
C	1	1
H	4	4
O	4	4

Since, the number of atoms of each element on both sides of the equation is same, the equation is balanced.


Hence, the balanced equation is: $CH_4 + 2O_2 \rightarrow CO_2 + 2H_2O$

Ex.4 Nitrogen and hydrogen combine to form ammonia.


Sol. Step-I: Writing the chemical equation in the word form:

Nitrogen + Hydrogen $\rightarrow$ Ammonia


Step-II: Writing the skeletal chemical equation:



Step-III: Balancing of atoms on both sides: There are 2 N-atoms on LHS and 1 N-atom on RHS. To balance N-atoms, multiply, NH_3 on RHS by 2 we get



Now, there are 6 H-atoms on RHS and 2 H-atoms on LHS. To balance H-atoms, multiply H_2 on LHS by 3, we get,



Step-IV: To check the correctness of the balanced equations:

Element	Number of atoms of on LHS	Number of Atoms on RHS
N	2	2
H	6	6

Since, the number of atoms of each element is equal on both sides of the equation, it is balanced.
Hence, the balanced equations is: $N_2 + 3H_2 \rightarrow 2NH_3$

COMPETITIVE LEVEL

♦ Information provided by a balanced chemical equations:

A skeletal equation provides only qualitative aspect of a chemical reaction. On the other hand, balanced chemical equation provides both qualitative and quantitative aspects.

Qualitative aspect of a chemical equation tells only which reactants are taking part in the reaction and which products are being formed in it.

e.g.: $C(s) + O_2(g) \rightarrow CO_2(g)$

The reaction suggests that solid carbon reacts with oxygen gas on heating to give carbon dioxide in gaseous form.

Quantitative aspect of a chemical equation provides the following information:

The stoichiometric ratio in which reactants are used and products are formed. The mole concept allows us to use the quantitative information available in a balanced chemical equation. The quantitative aspect, dealing with mass and volume relations among reactants and products is termed stoichiometry (derived from a Greek word meaning to measure an element).

Consider for example, the reaction represented by a balanced chemical equation:

Chemical Equation:

	$2\text{H}_2(\text{g})$	+	$\text{O}_2(\text{g})$	$\rightarrow$	$2\text{H}_2\text{O}(\text{g})$
Mole ratio:	2 mol		1 mol		2 mol
	or		or		or
Molecule:	$2 \times 6.023 \times 10^{23}$		6.023×10^{23}		$2 \times 6.023 \times 10^{23}$
ratio	or		or		or
	2 molecules		1 molecules		2 molecules
Mass ratio:	4g		32 g		36 g
Volume ratio:	2 vol		1 vol		2 vol

(Volume ratio is valid only for gaseous state at same P and T)

The given reaction suggest for the combination ratio of reactants and formation ratio of products in terms of:

- (a) **Mole ratio:** 2 mol H_2 reacts with 1 mol O_2 to form 2 mol H_2O vapours.
- (b) **Molecular ratio:** 2 molecule H_2 reacts with 1 molecule O_2 to form 2 molecules H_2O vapours.
- (c) **Mass ratio:** 4 g H_2 reacts with 32 g O_2 to form 36 g of H_2O vapours.
- (d) **Volume ratio:** In gaseous state 2 volume H_2 reacts with 1 volume O_2 to form 2 volume H_2O vapours at same conditions of Pressure and Temperature.

It is thus, evident that the coefficients in a balanced chemical equation can be interpreted as the relative number of moles, molecules or volume (if reactants are gases) involved in the reaction. These coefficients are called stoichiometric equivalent.

◆ **Limitations of a Chemical Equation:** It gives no information about the following:


- (i) The physical state of the reactants
- (ii) The concentration of the reactants.
- (iii) The conditions necessary to start and carry on the reaction e.g.: is any catalyst required.
- (iv) Is the reaction exothermic or endothermic, i.e., is heat evolved or absorbed during the reaction.
- (v) Reversibility or irreversibility of a reaction.


◆ **Modification Made for Improvement in Chemical Equations:**

Several modifications are introduced for making chemical equation more informative and more useful.

- (i) **The physical state of reactants and products:** The physical state of the reactants and products are specified by putting letters 's' (for the solid state), 'l' (for the liquid state) and 'g' (for the gaseous state) or 'aq' (for the aqueous solution) in parentheses just after the chemical formula of the reactant or the product which needs to be specified. Sometimes if a gas is evolved in a reaction it is shown by putting the symbol ($\uparrow$) of an arrow pointing upwards. Similarly if any precipitate is formed during the reaction, it is indicated by putting the symbol ($\downarrow$) of an arrow pointing downwards.


For example.






(ii) The concentration of an acid or a base:

The strength of an acid or a base solution is expressed by writing dil. for a dilute solution and conc. for a concentrated solution. For example,



(iii) Favourable conditions of the reactions: The conditions at which a given reaction is made are mentioned on the row between reactants and products.

For example,



This equation signifies that the nitrogen and hydrogen combines at 450°C and 200 atmospheric pressure in the presence of a catalyst iron (Fe) and promoter molybdenum (Mo).


(iv) Heat changes in the reaction:

Exchange of heat takes place during chemical reactions and therefore heat evolved or absorbed in a chemical reaction should be represented accordingly with their unit.

- Evolution of heat:

Reactants $\rightarrow$ Products + Heat


For example,



- Absorption of heat:

Reactants $\rightarrow$ Products – Heat

For example,





(v) Reversibility or irreversibility of a reaction:

This can be represented by use of different arrows in the equation. Double headed ($\rightleftharpoons$) or a double half headed ($\rightleftharpoons$) arrow is used for reversible while single headed ($\longrightarrow$) arrow is used for irreversible reactions.



- **Reversible reactions:**

For example,



- **Irreversible reactions:**

For example,



♦ **Limitations that are not rectified:**

(i) Rate of reaction:

To represent a slow and a fast reaction usually slow or fast is written on the arrow.

Time involved in the reactions.

(iii) Precautions to be taken during the reactions.

Types of Chemical Reactions

The chemical reactions can broadly be classified into following types:


- (1) Combination reaction
- (2) Decomposition reaction
- (3) Displacement reaction
- (4) Double displacement reaction
- (5) Redox reaction

1. Combination Reaction:

The chemical reaction involving combination of two or more reactants (elements or compounds) to form a single new product is called combination reaction.

Note: A reaction made to obtain a product by using two or more elements is usually referred as synthesis of that compound. Thus combination reaction is also known as synthesis reaction.


Combination reaction like other reactions also involves the exchange of energy i.e., heat, light electricity, pressure or catalyst e.g.:




Where, A and B are reactants while AB is the product.

♦ **Combination reaction between two elements-**


(i) Hydrogen burns in an atmosphere of oxygen, to form water only.



(ii) Combination of hydrogen and chlorine to form hydrogen chloride.




(iii) Combination of sulphur and oxygen to form sulphur dioxide.




♦ **Combination reaction between a compound and an element-**


(i) Combination of carbon monoxide and hydrogen to form methyl alcohol.



(ii) Combination of carbon dioxide with carbon to form carbon monoxide.




(iii) Combination of nitric oxide and oxygen to form nitrogen dioxide.




♦ **Combination reaction between two compounds-**

(i) Dissolution of sulphur trioxide in water to form sulphuric acid.




(ii) Dissolution of quick lime in water to form lime water.




The lime water solution, when applied to the walls of building for white washing, reacts slowly with $\text{CO}_2(\text{g})$ present in air and forms thin shining layer of CaCO_3 on the walls.

(iii) Combination of ammonia and chlorine to give ammonium chloride.




2. Decomposition Reaction:

The chemical reaction in which a compound is broken up into two or more simpler substances (elements or compounds) is called a decomposition reaction.



A single reactant AB decomposes to give simpler products like C, D etc.

Note: Decomposition reactions in which a compound breaks down into its constituent elements, is called analysis reactions.




Decomposition reaction involves absorption of energy e.g.: heat, light or electricity. Heat, light and electricity are different form of energy and causes breaking of bonds in the molecules of reactants to yield simpler products.

On the basis of form of energy which is supplied for decomposition, decomposition reactions can be classified as:

- ♦ **Thermal decomposition:** The decomposition reactants involving absorption of heat are called thermal decomposition reactions.

(i) Thermal decomposition of mercuric oxide: Mercuric oxide on heating strongly decomposes to give mercury and oxygen.



Experiment: Mercuric oxide a red coloured solid on heating decomposes to give shining droplets of mercury with the evolution of oxygen gas.




Fig. Heating of mercuric oxide

(ii) **Thermal decomposition of lead nitrate:**



Experiment: About 2 gram of lead nitrate is taken in a hard glass test tube, holding it by a test tube holder and heat is over a burner. Brown fumes of nitrogen dioxide gas are noticed at the mouth of the test tube and lead oxide is left behind in the test tube in the form of a yellow solid.




Fig. Thermal decomposition of lead nitrate

COMPETITIVE LEVEL

Oxidation Number

Definition: Oxidation number of atom in a molecule or a polyatomic ion is a hypothetical charge the atom would have if the electrons in each bond were located on the more electronegative atom.

Oxidation number may be integer, can be positive, negative or zero.

♦ **Calculating oxidation number:**

Following rules must be considered for calculating oxidation number.

Rule 1 : The oxidation number of each atom in an elemental molecule is always zero.
e.g.: Oxidation number of N, O, Cl, F, H, Na in N_2 , O_2 , Cl_2 , F_2 , H_2 , Na is zero.

Rule 2 : The sum of oxidation number of all atoms in a molecule is always zero.

Rule 3 : The sum of oxidation number of all atoms in an anion is equal to the total charge present on an ion.

Rule 4 : The more electronegative element in a molecule or ion has a negative oxidation number while the less electronegative has positive oxidation number.

Rule 5 : The oxidation number of H in most of its compounds is +1.

Rule 6 : The oxidation number of O in most of its compounds is -2.

Rule 7 : The oxidation number of F in all of its compounds is always -1.

Rule 8 : Oxidation number of alkali metals (Li, Na, K, Rb, Cs, Fr) is + 1.

Rule 9 : Oxidation number of alkaline earth metals (Be, Mg, Ca, Sr, Ba) is + 2.


Rule 10 : Oxidation number of halogens (Cl, Br, I, At) is usually -1.

Oxidation and Reductions in Day-to-Day Life

Redox changes are important reactions which influence our day-to-day life in several ways. Some of its examples such as burning of fuels, digestion of food in our body etc. are boon to mankind and are very helpful in sustaining life. On the other hand, some of its effects are quite damaging also e.g.: burning of fuels causing air pollution, rancidification of food, corrosion of metals etc.


◆ Useful Effects of Oxidation

1. **Combustion Reaction:** Kerosene, coal, charcoal, wood etc. burn in air and undergo combustion. Methane (CH_4) a major constituent of natural gas undergoes combustion in excess of oxygen upon heating.



Methane

Similarly, butane (C_4H_{10}) a main constituent of L.P.G. also undergoes combustion.



Butane

All combustion reactions are exothermic and redox reactions in nature and release heat energy. The human body may be regarded as a furnace or machine to which various food stuffs that we eat undergo combustion or oxidation. The heat energy evolved keeps our body working. Carbohydrates such as glucose, fructose, starch etc, are the major source of energy to the human body. They undergo combustion with the help of oxygen that we inhale to form carbon dioxide and water. For **Example.**



All combustion reactions are not accompanied by flame. Combustion is basically oxidation accompanied with release of energy.

EXERCISE-1

➤ Very Short Answer Type Questions

- Q.1** Write balanced equation for the following reaction:
Zinc carbonate_(s) $\longrightarrow$ Zinc oxide_(s)
+ Carbon dioxide_(g)
- Q.2** Give one example of an electrolytic decomposition.
- Q.3** What happens when sodium reacts with water?
- Q.4** Why is photosynthesis considered as an endothermic reaction?
- Q.5** Give an example of chemical reaction characterized the change in temperature.

➤ Short Answer Type Questions – Type I

- Q.6** What is a decomposition reaction? Give example.
- Q.7** X, Y and Z are three elements which undergo chemical reactions according to following equations.
 $X_2O_3 + 2Y \rightarrow Y_2O_3 + 2X$
 $3ZSO_4 + 2Y \rightarrow Y_2(SO_4)_3 + 3Z$
 $3ZO + 2X \rightarrow X_2O_3 + 3Z$
Answer the following equations:
(i) Which element is the most reactive?
(ii) Which element is the least reactive?
- Q.8** Potassium chlorate (KClO₃) on heating forms potassium chloride and oxygen. Write a balanced equation for this reaction.
- Q.9** On the basis of the following chemical equations, find out which is the least reactive metal amongst iron, copper and zinc?
(i) $FeSO_4(aq) + Zn(s) \rightarrow ZnSO_4(aq) + Fe(s)$
(ii) $CuSO_4(aq) + Fe(s) \rightarrow FeSO_4(aq) + Cu(s)$

- Q.10** What type of chemical equation are the following equations:

- (i) $A + BC \longrightarrow AC + B$
(ii) $A + B \longrightarrow AB$
(iii) $AB \longrightarrow A + B$
(iv) $AB + CD \longrightarrow AD + CB$

➤ Short Answer Type Questions – Type II

- Q.11** How do we come to know that a chemical reaction has taken place?
- Q.12** In the following situations, a reaction may or may not take place, If it does, write the balanced equations illustrating the reaction. Assume all involve aqueous solutions.
(i) Some iron nails are placed in a CuCl₂ solutions.
(ii) Silver coins are dropped in a hydrochloric acid solution.
(iii) A copper wire is placed in a Pb(NO₃)₂ solution.
- Q.13** Consider the following chemical equations:
(i) $CuO(s) + H_2(g) \rightarrow Cu(s) + H_2O(g)$
(ii) $Fe_2O_3(s) + 3CO(g) \rightarrow 2Fe(l) + 3CO_2(g)$
Identify the following in these equations, giving reasons:
(a) The substance getting oxidised.
(b) The substance getting reduced
(c) The oxidising agent
(d) The reducing agent
- Q.14** Among the following displacement reactions which one will take place and which one will not occur and why?
(i) $MgSO_4(aq) + Zn(s) \rightarrow ZnSO_4(aq) + Mg(s)$
(ii) $CuSO_4(aq) + Fe(s) \rightarrow FeSO_4(aq) + Cu(s)$
- Q.15** What is an oxidation reaction? Give an example of oxidation reaction. Is oxidation an exothermic or an endothermic reaction.

- Q.16** What happens when iron nails are put in copper sulphate solution?
 (i) Write the equation for the reaction that takes place.
 (ii) Name the type of reaction involved.
- Q.17** Why does stale food give a bad taste and bad smell?
- Q.18** Why do silver, gold and platinum not corrode in moist air?
- Q.19** What are the different types of combination reactions?
- Q.20** Why cannot we stir silver nitrate solution with copper spoon?

➤ Long Answer Type Questions

- Q.21** Write any two observations in an activity which may suggest that a chemical reaction has taken place. Give an example in support of your answer.
- Q.22** Balance the following chemical equations:
 (i) $S(s) + H_2SO_4(aq) \longrightarrow H_2O(l) + SO_2(g)$
 (ii) $S(s) + HNO_3(aq) \longrightarrow$
 $H_2SO_4(aq) + NO_2(g) + H_2O(l)$
 (iii) $Fe_2O_3(s) + CO(g) \longrightarrow Fe(l) + CO_2(g)$
 (iv) $KMnO_4(aq) + HCl(aq) \longrightarrow$
 $KCl(aq) + MnCl_2(aq) + Cl_2(g) + H_2O(l)$
 (v) $MnO_2(s) + HCl(aq) \longrightarrow$
 $MnCl_2(aq) + H_2O(l) + Cl_2(g)$
- Q.23** Name the substance oxidized and substance reduced in the following reactions:
 (i) $MnO_2 + 4HCl \rightarrow MnCl_2 + Cl_2 + 2H_2O$
- (ii) $2H_2S + SO_2 \rightarrow 2H_2O + 3S$
 (iii) $2PbO + C \rightarrow 2Pb + CO_2$
 (iv) $PbS + 4H_2O_2 \rightarrow Pb + CO_2$
- Q.24** Explain the effects of oxidation reactions in everyday life.
- Q.25** When a compound of lead is heated, brown coloured fumes evolve from it.
 (i) Name the compound.
 (ii) Write the balanced chemical equation.
 (iii) Write the chemical name of brown coloured fumes.
 (iv) Name the type of reaction.
 (v) Is this reaction an exothermic or endothermic ?

➤ Practical & Value Based Type Questions

- Q.26** Solutions of ferrous sulphate, zinc sulphate, copper sulphate and aluminium sulphate were separately taken in four test-tubes and some iron nails were placed in each of the solution. What will you observe after few seconds?
- Q.27** On keeping iron nails in blue coloured copper sulphate solution, it is observed that the colour of the solution turns light green after some time. Give reason for this colour change. Name the type of this reaction.

EXERCISE-2

- Q.1** Which of the following changes represents a physical change?
 (A) Conversion of sulphur to sulphur dioxide
 (B) Heating of a metal knife
 (C) Combustion of methane
 (D) Hydrolysis of an ester
- Q.2** Which of the following are exothermic processes?
 (i) Reaction of water with quick lime
 (ii) Dilution of an acid
 (iii) Evaporation of water
 (iv) Sublimation of camphor (crystals)
 (A) (i) and (ii)
 (B) (ii) and (iii)
 (C) (i) and (iv)
 (D) (iii) and (iv)
- Q.3** Which of the following is not an endothermic process?
 (A) Boiling of water
 (B) Dissolution of salts in water
 (C) Dissolution of conc. H_2SO_4 in water
 (D) Evaporation of water
- Q.4** If in the reaction $\text{N}_2(\text{g}) + \text{O}_2(\text{g}) \longrightarrow 2\text{NO}(\text{g})$ the energy required to break nitrogen-nitrogen and oxygen-oxygen bonds is more than twice the energy released during bond formation between nitrogen and oxygen atoms, the reaction is a/an -
 (A) Endothermic reaction
 (B) Double displacement reaction
 (C) Exothermic reaction
 (D) Single displacement reaction
- Q.5** In an endothermic reaction -
 (A) The energy content of products is less than heat content of reactants
 (B) The energy content of products is greater than that of reactants
 (C) Heat is released
 (D) Heat is neither absorbed nor released
- Q.6** Three beakers labeled as A, B and C each containing 25 ml of water were taken. A small amount of NaOH, anhydrous CuSO_4 and NaCl were added to the beakers A, B and C respectively. It was observed that there was an increase in the temperature of the solutions contained in beakers A and B, whereas, in case of beaker C, the temperature of the solution falls. Which one of the following statement (s) is (are) correct?
 (i) In beakers A and B, exothermic process has occurred.
 (ii) In beakers A and B, endothermic process has occurred.
 (iii) In beaker C exothermic process has occurred.
 (iv) In beaker C endothermic process has occurred.
 (A) (i) only (B) (ii) only
 (C) (i) and (iv) (D) (ii) and (iii)
- Q.7** Complete the following statement by choosing correct options for x and y. "During the process of respiration, glucose combines with oxygen in the cells of our body and 'x' a large amount of energy. Hence, respiration is an 'y' process."
 (A)

x	y
releases	endothermic

 (B)

x	y
absorbes	endothermic

 (C)


x	y
releases	exothermic

 (D)

x	y
absorbes	exothermic
- Q.8** A balanced chemical equation is in accordance with -
 (A) Avogadro's law
 (B) Dalton's atomic theory
 (C) The law of conservation of mass
 (D) The law of constant proportion

- Q.9** Which of the following chemical equations is an unbalanced one?
- (A) $2\text{NaHCO}_3 \longrightarrow \text{Na}_2\text{CO}_3 + \text{H}_2\text{O} + \text{CO}_2$
- (B) $2\text{C}_4\text{H}_{10} + 12\text{O}_2 \longrightarrow 8\text{CO}_2 + 10\text{H}_2\text{O}$
- (C) $2\text{Al} + 6\text{H}_2\text{O} \longrightarrow 2\text{Al}(\text{OH})_3 + 3\text{H}_2$
- (D) $4\text{NH}_3 + 5\text{O}_2 \longrightarrow 4\text{NO} + 6\text{H}_2\text{O}$

- Q.10** Which of the following statements is not true for the chemical equation?




- (A) 2 mol of N_2O_5 on dissociation gives 4 mol of NO_2 and 1 mol of O_2
- (B) 1 mol of N_2O_5 on dissociation gives 2 mol of NO_2 and 0.5 mol of O_2
- (C) 2 g of N_2O_5 on dissociation gives 4 g of NO_2 and 1 g of O_2
- (D) 216 g of N_2O_5 on dissociation gives 184 g of NO_2 and 32 g of O_2

- Q.11** Electrolysis of water is a decomposition reaction. The mole ratio of hydrogen and oxygen gases liberated during electrolysis of water is -

- (A) 1: 1 (B) 2: 1
(C) 4: 1 (D) 1: 2

- Q.12** Take a long magnesium ribbon. Heat it over a flame when it burns with a bright light and is converted into white powder of MgO . It is called a -
- (A) combination reaction
(B) displacement reaction
(C) decomposition reaction
(D) thermal decomposition

- Q.13** Which of the following reactions should be called 'combination' or 'synthesis' reaction?
- (i) $\text{C}(\text{s}) + \text{O}_2(\text{g}) \longrightarrow \text{CO}_2(\text{g})$
- (ii) $\text{CaO}(\text{s}) + \text{CO}_2(\text{g}) \longrightarrow \text{CaCO}_3(\text{s})$







- (A) Reaction (i) only
(B) Reactions (i) and (ii) only
(C) Reactions (i), (ii) and (iii) only
(D) All reactions – (i) to (iv)

- Q.14** Take about 1.0 g CaCO_3 in a test tube. Heat it over a flame, when a colourless gas comes out. The reaction is called a -


- (A) decomposition reaction
(B) displacement reaction
(C) double decomposition reaction
(D) double displacement reaction

- Q.15** Which of the following reactions should be called 'decomposition' reaction?



- (A) Reaction (i) only
(B) Reactions (i) and (ii) only
(C) Reactions (i), (ii) and (iii) only
(D) All reactions – (i) to (iv)

- Q.16** The following reaction is used for the preparation of oxygen gas in the laboratory:



Which of the following statement(s) is/are correct about the reaction?

- (A) It is a decomposition and endothermic in nature.
(B) It is a combination reaction.
(C) It is a decomposition reaction and accompanied by release of heat
(D) It is a photochemical decomposition reaction and exothermic in nature

EXERCISE-3


- Q.1** The reaction between aqueous solutions of sodium chloride and silver nitrate is –
[Raj. NTSE Stage-I/05]
(A) displacement reaction
(B) synthesis reaction
(C) double displacement reaction
(D) analysis reaction
- Q.2** $\text{CH}_4 + \text{Cl}_2 \rightarrow \text{CH}_3\text{Cl} + \text{HCl}$ reaction is an example of - [Raj. NTSE Stage-I/06]
(A) Synthetic (B) Analytical
(C) Displacement (D) Neutralisation
- Q.3** A Brown and bright element “x” when heated in presence of air turns into black substance “y”. If Hydrogen gas is passed over this heating material again “x” is obtained. “x” and “y” are –
[Raj. NTSE Stage-I/13]
(A) Cu & CuO (B) S & SO_2
(C) C & CO_2 (D) Na & NaH
- Q.4** $\text{H}_2\text{S}(\text{g}) + \text{Cl}_2(\text{g}) \rightarrow 2\text{HCl}(\text{g}) + \text{S}(\text{s})$
The reaction is interpreted as:
[Delhi. NTSE Stage-I/13]
(A) H_2S is getting oxidised and Cl_2 is getting reduced
(B) H_2S is getting reduced and Cl_2 is getting oxidised
(C) Only H_2S is oxidised
(D) Both H_2S and Cl_2 are reduced
- Q.5** What is the oxidation number of sulphur in peroxy mono sulphuric acid (H_2SO_5)?
[Haryana. NTSE Stage-I/13]
(A) 8 (B) 6 (C) 5 (D) 4
- Q.6** Oxidation is defined as:
[M.P. NTSE Stage-I/13]
(A) loss of electron (B) gain of electron
(C) loss of proton (D) gain of proton
- Q.7** From the following metals whose nitrate produces NO_2 gas on heating?
[West Bengal NTSE Stage-I/13]
(A) Na (B) K
(C) Pb (D) None of these
- Q.8** The equation given below indicates
 $\text{NaCl} + \text{AgNO}_3 \rightarrow \text{AgCl} + \text{NaNO}_3$;
[Karnataka. NTSE Stage-I/13]
(A) Chemical decomposition
(B) Chemical combination
(C) Chemical displacement
(D) Chemical double displacement
- Q.9** Displacement reaction is:
[Raj. NTSE Stage-I/14]
(A) $\text{CaO}(\text{s}) + \text{H}_2\text{O}(\text{l}) \rightarrow \text{Ca}(\text{OH})_2(\text{aq})$
(B) $\text{Pb}(\text{s}) + \text{CuCl}_2(\text{aq}) \rightarrow \text{PbCl}_2(\text{aq}) + \text{Cu}(\text{s})$
(C) $\text{MnO}_2(\text{s}) + 4\text{HCl}(\text{l}) \rightarrow \text{MnCl}_2(\text{s}) + 2\text{H}_2\text{O}(\text{l}) + \text{Cl}_2(\text{g})$
(D) $\text{C}_6\text{H}_{12}\text{O}_6 + 6\text{O}_2 \rightarrow 6\text{CO}_2 + 6\text{H}_2\text{O}$
- Q.10** What is the instrument called for water electrolysis process?
[Gujrat NTSE Stage-I/14]
(A) Voltmeter (B) Voltmeter
(C) Hydrometer (D) Lectometer
- Q.11** Which of the following is endothermic reaction? [Haryana NTSE Stage-I/14]
(A) $\text{C}(\text{s}) + \text{O}_2(\text{g}) \longrightarrow \text{CO}_2(\text{g})$
(B) $\text{N}_2(\text{g}) + \text{O}_2(\text{g}) \longrightarrow 2\text{NO}(\text{g})$
(C) $2\text{H}_2(\text{g}) + \text{O}_2(\text{g}) \longrightarrow 2\text{H}_2\text{O}(\text{l})$
(D) $2\text{CH}_3\text{OH}(\text{l}) + 3\text{O}_2(\text{g}) \rightarrow 2\text{CO}_2(\text{g}) + 4\text{H}_2\text{O}(\text{l})$
- Q.12** In which of the following reactions H_2O_2 acts as a reducing agent:
[Bihar NTSE Stage-I/15 & JEE main/14]
(a) $\text{H}_2\text{O}_2 + 2\text{H}^+ + 2\text{e}^- \rightarrow 2\text{H}_2\text{O}$
(b) $\text{H}_2\text{O}_2 - 2\text{e}^- \rightarrow \text{O}_2 + 2\text{H}^+$
(c) $\text{H}_2\text{O}_2 + 2\text{e}^- \rightarrow 2\text{OH}^-$
(d) $\text{H}_2\text{O}_2 + 2\text{OH}^- - 2\text{e}^- \rightarrow \text{O}_2 + 2\text{H}_2\text{O}$
(A) (a),(c) (B) (b),(d)
(C) (a),(b) (D) (c), (d)

Q.13 Precipitate formation during chemical reaction is indicated by this arrow:

[Maharashtra_NTSE Stage-I/15]

(A) $\uparrow$ (B) $\rightarrow$ (C) $\downarrow$ (D) $\leftarrow$

Q.14 $\text{BaCl}_{2(aq)} + \text{Na}_2\text{SO}_{4(aq)} \longrightarrow$



[Maharashtra_NTSE Stage-I/15]

The types of reaction are:

(a) Displacement

(b) Precipitation

(c) Combination

(d) Double displacement


(A) (a) & (c)

(B) (a), (b), & (c)

(C) (b) & (c)

(D) (b) & (d)

Q.15 The chemical reaction



is an example of - [Raj._NTSE Stage-I/15]


(A) Neutralization

(B) Double displacement

(C) Neutralization and double displacement

(D) Combination

Q.16 Which type of catalyst is glycerol in the following reaction ? [Raj/NTSE/18/Stage-I/]



[Raj._NTSE Stage-I/18]

(A) Positive catalyst

(B) Negative catalyst

(C) Biocatalyst

(D) Autocatalyst

ANSWER KEY

EXERCISE - 2

Ques.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Ans.	B	A	C	A	A	C	C	C	B	C	B	A	C	A	D
Ques.	16														
Ans.	A														


EXERCISE - 3

Ques.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Ans.	C	C	A	A	A	A	C	D	B	A	B	B	C	D	C
Ques.	16														
Ans.	B														


NUTRITION

Chapter Outline


- ❖ Nutrition
- ❖ Modes of nutrition
- ❖ Nutrition in plants
- ❖ Nutrition in animals
- ❖ Digestive system of humans




Digestive System



Nutrition of Amoeba



Plant Showing Photosynthesis



NUTRITION

Nutrition

The process of intake of nutrients & its utilization is called nutrition.

◆ Nutrient :


The different component of food that have distinct functions like

- (i) providing energy
- (ii) providing materials for body building
- (iii) maintenance & regulation of metabolism are called nutrient.

For Example –

- (i) Proteins - Egg, pulses
- (ii) Minerals, Vitamin - Green leafy vegetable
- (iii) Carbohydrates - Wheat
- (iv) Fats - Butter, Oils

Modes of Nutrition



◆ Autotrophic Nutrition : (Gk. *autos*-self, *trophe*-nourishment)

It is a mode of nutrition in which organisms are able to build up their own organic food from inorganic raw materials with the help of energy. The organism performing autotrophic nutrition are called autotrophs.

◆ Heterotrophic Nutrition :

It is a mode of nutrition in which the organisms obtain readymade organic food from outside sources. The organisms that depend upon outside sources for obtaining organic nutrients are called heterotrophs. Heterotrophic nutrition is of three types - saprophytic, parasitic and holozoic.

Ex.1 What kind of nutrition occurs in yeast.

Ans. Saprophytic nutrition

COMPETITIVE LEVEL

- Fats provide the most energy followed by carbohydrates and proteins respectively.
- Proteins are the building blocks of body.
- **Chemoautotrophs** : These organisms use chemical energy for the synthesis of food.

E.g. Purple sulphur bacteria

- **Photoautotrophs** : These organism use light energy for the synthesis of food.

E.g. Green plants


- **Saprophytic or Saprotrophic Nutrition** : It is a mode of heterotrophic nutrition in which food is obtained from organic remains like dead organisms, excreta, fallen leaves, broken twigs, food articles, etc.
- **Parasitic Nutrition** : It is a mode of hetrotrophic nutrition in which a living organism flourishes by obtaining food from another living organism.
- **Holozoic Nutrition** : It is a mode of heterotrophic nutrition which involves intake of solid pieces of food.

Nutrition in Plants

◆ Photosynthesis :

Photosynthesis is the primary mode of food production in green plant. " The process by which green plants synthesize food from simple substances i.e. carbon dioxide and water in the presence of light and chlorophyll is called **Photosynthesis**"

The process of photosynthesis can be represented in the form of chemical reaction, as given below :



◆ Essential of photosynthesis :

- (A) **Sunlight** : For plants sun is the basic source of radiant energy.
- (B) **Chlorophyll** : These are the green pigments present in chloroplast. They are found in green leaves in the maximum amount as well as in other green aerial parts of plant.




Figure : Chloroplast

- (C) **Water** : Plant's roots absorb water from the soil by the process of osmosis (endosmosis). This water is transported to leaves by a special type of tissue called as xylem.

(D) **Carbon Dioxide** : Terrestrial plants obtain carbon dioxide from the atmosphere through the small openings present on leaves called as stomata. 'Stomata' are the small pores present on the surface.

◆ Opening and Closing Stomata :

The opening and closing of stomata depend upon the turgid or flaccid state of the guard cells. When guard cells are in turgid state the stomatal aperture opens and when guard cells are in flaccid state the stomatal aperture closes.

The inner wall of guard cells (towards pore) is thick and outer wall (towards other epidermal cells) is thin. When the turgor pressure of the guard cells is increased the outer thinner wall of the guard cell is pushed out (towards the periphery) due to which a tension is created on the inner thicker wall thus pulling the inner thicker wall towards the periphery thus leading to the opening of stomatal aperture.

On the contrary when the guard cells are in a flaccid state the outer thinner wall of guard cells returns to original position (moves towards pore) due to which tension on the inner wall is released which also returns to its original position and stomatal aperture gets closed again.




Figure : Stomata

COMPETITIVE LEVEL

- Plants utilize the light in the visible region of solar spectrum (electromagnetic spectrum) which comes under the range of 390 nm – 780 nm wavelength.
- Photosynthetically Active Radiation or PAR refers to radiation with wavelengths between 400 & 700 nm.
- 50% of incident solar radiation is PAR. Plant absorbs only 2-10% of PAR.
- Maximum photosynthesis takes place in white light followed by red and blue light.
- Minimum photosynthesis takes place in green light.
- RuBisCO is the most abundant enzyme.
- Plants utilize the light in the visible region of solar spectra (electromagnetic spectrum) which comes under the range of 390 nm – 780 nm wavelength
- Maximum photosynthesis occurs in red region
- There are six different types of chlorophyll : chl a, b, c, d, e and bacteriochlorophyll. Amongst them chlorophyll a and chlorophyll b are the most commonly occurring chlorophyll.

◆ Mechanism of Photosynthesis :

The following events occur in photosynthesis

- (i) Absorption of light energy by chlorophyll.
- (ii) Conversion of light energy to chemical energy and splitting of water molecules into hydrogen and oxygen.
- (iii) Reduction of carbon dioxide to carbohydrates.

COMPETITIVE LEVEL

◆ Photosynthesis can also be Divided into Two Steps :

(A) Light Reaction :


- It is also called Hill Reaction. As it was discovered by Robert Hill.
- Site - grana of thylakoids
- It is named as light reaction as it occurs only in presence of light.

Major Step :

- (i) **Photoexcitation of chlorophyll molecule** : During this process chlorophyll molecule receives sunlight in the form of small energy bundles called as **photons** and becomes excited to higher energy level.
- (ii) **Photolysis** : It is also called as photooxidation of water, this takes place in presence of Mn^{+2} and Cl^{-} ions.



O_2 is liberated as by product and H^{+} ions are used for reduction of NADP



- (iii) **Photophosphorylation** : During this process ATP are produced. It takes place in **quantasomes** / **photosystem**. Inorganic phosphate is required to convert ADP to ATP.

(B) Dark Reaction :

- This reaction is not dependent on light. It is also known as Calvin – Benson Cycle or C_3 cycle as first stable product is phosphoglyceric acid (PGA) a 3 carbon compound.
- **Site** : Stroma of chloroplast

Major Step :

- (i) **Carboxylation** : In this CO_2 is captured by CO_2 acceptors like RUBP (C_3 Plants) PEP (C_4 Plants) with the help of **carboxylase enzyme** i.e. **RuBisCO & PEPCo** respectively. The first stable compound of C_3 cycle is a three carbon containing molecule **phosphoglyceric acid** so it is called as C_3 cycle while in C_4 cycle it is four carbon containing **oxalo acetic acid**.
- (ii) **Synthesis of glucose** : In this phase captured CO_2 is assimilated into glucose in the presence of phosphatase and isomerase enzyme.
- (iii) **Regeneration of RUBP.**

Ex.2 Why are plants green?

Ans. Plants are green due to presence of the green pigment called chlorophyll.

Ex.3 “Plants prepare their own food and are called autotrophs”. What is the name of this process ? Why is it called so?

Ans. The process of production of food by plants is called photosynthesis. Photo means light and synthesis means manufacturing, as manufacturing of glucose occurs in the presence of light it is named as ‘photosynthesis’.

Ex.4 Oxygen released during photosynthesis is the outcome of which steps of this process?

Ans. The energy trapped by the chlorophyll molecules is used to breakdown water and released oxygen as a product. This process is called photolysis of water.

COMPETITIVE LEVEL

◆ Crassulacean Acid Metabolism (CAM) :

- Certain plants, especially succulents which grow under extremely xeric (dry) condition, fix atmospheric CO₂ in dark.
- Since the process was first observed in the plants belonging to family crassulaceae (eg. Bryophyllum, kalanchoe etc.) It was termed crassulacean acid metabolism (CAM).
- The most characteristic feature of these plants is that their stomata remain open at night (in dark) but closed during the day (in light).
- Thus, CAM is a kind of adaptation in succulents to carry out photosynthesis without much loss of water.

Nutrition in Animals

◆ Nutrition in Amoeba :

Protozoa carry out holozoic nutrition through intracellular digestion.

- Ingestion :** Some protozoa can ingest food particle from any point on the surface (e.g., *Amoeba*) while others have fixed points for the same (e.g., *Paramecium*). Protozoans like *Amoeba* capture food with the help of temporary finger-like processes called **pseudopodia**. Protozoans like (*Paramecium* have small hair-like processes called cilia.) Beating of cilia creates current in water that pushes food particle through cytostome or cell mouth. The process of ingestion of solid food particle by a cell or unicellular organism is called **phagocytosis**.
- Digestion :** In amoeba, food is digested in the food vacuole by digestive enzymes. These enzymes break down the food into small and soluble molecules by chemical reactions.
- Absorption :** The digested food is directly absorbed into the cytoplasm by diffusion. After absorption the food vacuole disappears.
- Assimilation :** The absorbed food is used to obtain energy which leads to growth and reproduction of Amoeba.

Digestive System of Human

Digestive system is a group of organs & associated digestive glands that take part in ingestion, digestive absorption of food & egestion of undigested matter. Digestive organs form a continuous canal called alimentary canal. Alimentary Canal in man is 9 metres long & consists of the following parts.





Figure : Digestive System

Ex.5 Why is digestion in Amoeba said to be intracellular ?

Ans. Intracellular means within a cell. Amoeba is a single-celled animals. All the life activities have to be performed within a single cell. Amoeba engulfs its food from the surrounding environment. Digestion, absorption, and assimilation occur inside the cell, hence it is said to exhibit intracellular mode of digestion.

◆ Associated Glands :

(A) Salivary glands (B) Gastric glands (C) Liver (D) Pancreas

COMPETITIVE LEVEL

- Salivary glands are of three types :
 - (i) Parotid gland
 - (ii) Sub mandibular glands,
 - (iii) Sub lingual glands
- Gastric glands are of three types :
 - (i) Cardiac gland

(ii) Pyloric gland

(iii) Fundic gland

- Liver is a bilobed structure and is the largest gland present in the human body.
- Liver is the only organ that possess capacity to regenerate in human.

◆ **Mouth :**

Transverse slit like aperture which is bounded by lips.

◆ **Oral Cavity :**

It has teeth, tongue & palate.

(A) Teeth :




Figure : Tooth

- The food taken inside oral cavity is masticated i.e. mechanically broken into smaller particles before being swallowed by the help of teeth.
- Man possesses teeth on both the jaws, there are 32 teeth of four different types, namely incisors, canines, premolars & molars.
- There are four different types of teeth in humans (Heterodont).

Incisors : For cutting

Canines : For tearing

Premolars : For grinding

Molars : For grinding

◆ **Dental Caries :**

Dental caries or tooth decay causes gradual softening of enamel and dentine. It begins when bacteria acting on sugars produce acids that softens or demineralises the enamel. Masses of bacterial cells together with food particles stick to the teeth to form dental plaque. Saliva cannot reach the tooth surface to neutralize the acids as plaque covers the teeth. Brushing the teeth removes the plaque before the bacteria produce acids. If untreated, microorganisms may invade the pulp, causing inflammation and infection.

Note : Brushing teeth properly avoids this situation.

COMPETITIVE LEVEL

◆ Dental Formula :

(i) Milk teeth / Primary

(ii) Permanent teeth

$$\frac{\text{No. of teeth in half part of upper jaw}}{\text{No. of teeth in half part of lower jaw}} \times 2 = x$$

- In Child $\rightarrow I \frac{2}{2}, C \frac{1}{1}, Pm \frac{0}{0}, M \frac{2}{2} = \frac{5}{5} \times 2 = \text{Total 20 teeth}$
- In Man $\rightarrow I \frac{2}{2}, C \frac{1}{1}, Pm \frac{2}{2}, M \frac{3}{3} = \frac{8}{8} \times 2 = \text{Total 32 teeth}$
- Jaws present in buccal cavity of humans are provided with four different types of teeth this is called **Heterodont**.
- Thecodont means “Socket tooth”, means that thecodont teeth are set in sockets of jaw bones.
- **Dental plaque** is a biofilm or mass of bacteria that grows on surfaces within the mouth.
- **Diastema** : It is a space or gap between two teeth.
- **Enamel** : It is the hardest substance of human body. It covers the crown portion of the tooth.
- **Dentin** : It is the hard dense bony tissue forming bulk of teeth & it is second hardest tissue in body after enamel.
- **Periodontium** : It refers to the specialized tissues that surround & support the teeth, maintaining them in their respective bones.

(B) Tongue

- The fleshy muscular organ in the mouth of a mammal, used for tasting, licking, swallowing and (in humans) articulating speech.
- It forms the floor of the mouth and bears taste buds.

◆ Pharynx :

- It is short conical region that lies after the oral cavity.
- It is a common passage to air and food.

◆ Oesophagus :

- It is a long narrow muscular tube which leads to the stomach. No digestive gland are present.
- It is also called as gullet or food pipe.
- **Peristalsis** is a series of contraction and relaxation of muscles of alimentary canal that pushes the food downward.

COMPETITIVE LEVEL

- Cooked starch $\xrightarrow[\text{Ptyalin}]{\text{Amylase}}$ Maltose
- Reverse peristalsis or retroperistalsis is the reverse of peristalsis, usually occur before vomiting.

◆ Stomach :

- It lies below the diaphragm on the left side of abdominal cavity.
- It is J-shaped. Its muscular walls help in mixing of food with the juices it produces.
- The entry and exit of food to and from the stomach is regulated by sphincter muscles.
- Food is churned in the stomach for about three hours.
- HCl provides an acidic medium to the food, and kills microorganisms present in the food.
- Mucus secreted by the stomach wall protects its wall from its own secretion of HCl.
- Peptic ulcers are sores or ulcers in the lining of the stomach, lower oesophagus, or small intestine, usually as a result of inflammation caused by the bacteria.
- Salivary amylase becomes inactive in stomach due to acidic pH.
- $\text{Pepsinogen} \xrightarrow{\text{HCl}} \text{Pepsin}$
 $\text{Protein} \xrightarrow{\text{Pepsin}} \text{Proteases}$
- Gastric lipases partially digest fats
- $\text{Prorenin} \xrightarrow{\text{HCl}} \text{Rennin}$
- $\text{Casein} \xrightarrow{\text{Rennin}} \text{Paracasein}$

◆ Small Intestine :

- It is a highly convoluted tube, and the site of complete digestion of food.
- Last part of small intestine is folded to form villi, which absorb the products of digestion
- It receives three kinds of juices :
 - (A) Pancreatic juice from pancreas
 - (B) Bile juice from liver
 - (C) Intestinal juice from the walls of intestine
- Liver secretes “Bile” which provides an alkaline medium and emulsifies the fat molecules.
- Duodenal wall secretes enterokinase which activates the trypsin
 $\text{Trypsinogen} \xrightarrow{\text{Enterokinase}} \text{Trypsin}$
 $\text{Chymotrypsinogen} \xrightarrow{\text{Trypsin}} \text{Chymotrypsin}$
 $\text{Polypeptides} \xrightarrow{\text{Chymotrypsin}} \text{Peptide fragments}$
- Carbohydrates are converted to disaccharides by pancreatic amylase and maltase converts them to glucose.
- Pancreatic lipase converts emulsified fats to fatty acids and glycerols.
- $\text{Fats} \xrightarrow[\text{Emulsify}]{\text{Bile}} \text{Fat globules} \xrightarrow{\text{Lipase}} \text{Glycerol} + \text{Fatty acid}$
- It is the longest in herbivores and smallest in carnivores as cellulose has to be digested in herbivores.

Ex.6 What is the function of digestive enzymes ?

Ans. The major constituents of the diet are relatively complex, such as carbohydrate, protein, (at, etc. which cannot be absorbed unless they are broken down into simple compounds. The function of digestive enzymes is to help in breaking down of complex food materials into simpler compounds which can be readily used by animaler through absorption and assimilation. As such, digestive enzymes help in converting proteins into amino acids, fats into fatty acids, and glycerols and polysaccharides into monosaccharides.

Ex.7 Write any two regions of alimentary canal in which no digestion occur?

Ans. Oesophagus, Large intestine.

Ex.8 What is the importance of Sphincters?

Ans. It regulates the exit of food in small amounts.

Ex.9 What is the role of saliva in the digestion of food?

Ans. Saliva contains salivary amylase and is released in our mouth. It breaks down starch into sugar (complex carbohydrates into simpler ones).

Ex.10 Why do herbivores have longer where as carnivores have shorter small intestine ?

Ans. Herbivores eat grass which has maximum cellulose content. It needs greater time for digestion. Carnivores eat meat which is easier to digest. Therefore, herbivores need longer small intestine and carnivores need shorter small intestine.

COMPETITIVE LEVEL

- The duodenum part of small intestine receives secretions from Liver and Pancreas through a common duct called the “Hepato-Pancreatic Duct”.
- Differentiated into 3 regions, viz. **Duodenum** which is the first part of small intestine & is curved C-shaped; **Jejunum**, comparatively longer & more coiled and **Ileum**,

◆ Large Intestine :

- It is much shorter & wider than small intestine
- No digestion takes place in large intestine, only absorption of water takes place.
- In herbivores like horse, rabbit digestion of cellulose takes place in caecum.

◆ Absorption :

- Some simple sugars are absorbed in the mouth.
- In the small intestine (ileum) absorption of all digested materials takes place through villi.
- Excess water is absorbed by the large intestine.

EXERCISE-1

➤ Very Short Answer Type Questions

- Q.1 Define villi.
- Q.2 What are heterotrophs ?
- Q.3 Which types of organisms are called consumers ?
- Q.4 What is the role of oesophagus ?
- Q.5 Define saprophyte.
- Q.6 What are the functions of liver in humans ?
- Q.7 Which type of animal is called omnivore?
- Q.8 Define digestion.
- Q.9 What is ingestion?
- Q.10 What is dental plaque?

➤ Short Answer Type Questions – Type I

- Q.11 What type of digestion occurs in *Paramoecium*?
- Q.12 How do saprophytic organisms obtain their nourishment?
- Q.13 Why chlorophyll is needed for photosynthesis?
- Q.14 Name a digestive juice that has no enzymes. What is the role of this juice?
- Q.15 Name the various parts of large intestine. What is the role of large intestine?

➤ Short Answer Type Questions – Type II

- Q.16 How does nutrition occur in *Amoeba*?
- Q.17 Differentiate between autotrophic and heterotrophic nutrition.
- Q.18 Distinguish saprophytes from parasites.
- Q.19 Differentiate between photosynthetic and holozoic nutrition.
- Q.20 What is the action of hydrochloric acid of gastric juice ?

➤ Long Answer Type Questions

- Q.21 Explain the mechanism of photosynthesis.
- Q.22 Describe the various types of heterotrophic nutrition.
- Q.23 Briefly describe the digestive system of humans.
- Q.24 What happens to food in the small intestine ?
- Q.25 What is the physical digestion of food ? Where does it occur ? What is its importance ?

➤ Practical & Value Based Questions

- Q.26 Om, Rohit and Kishore always remain in a hurry. One day, during lunch hour they all quickly gulped food and went out to play. (i). Om suddenly developed stomach ache while playing. What according to you might have gone wrong with him.
- Q.27 One day Mohan had a severe toothache. His father took Mohan to a dentist. The dentist examined all the teeth of Mohan very carefully and said that he had tiny holes in his two teeth. He also told Mohan that all his teeth were covered with a sticky, yellowish layer. The dentist performed a certain procedure on his two teeth having tiny holes and also gave him some medicines. Mohan's toothache disappeared gradually.
- What are the tiny holes in the teeth known as ?
 - How are the tiny holes formed in the teeth ?
 - What kind of procedure was performed by dentist on Mohan's two teeth ?
 - What is the sticky, yellowish layer on Mohan's teeth known as ?
 - How is the sticky, yellowish layer formed on the teeth ?
 - What advice will you give to Mohan to avoid such dental problems in future ?

EXERCISE-2

- Q.1** CO₂ and O₂ balance in atmosphere is due to
 (A) Photorespiration
 (B) Photosynthesis
 (C) Respiration
 (D) Transpiration
- Q.2** During photosynthesis the oxygen in glucose comes from
 (A) Water
 (B) Carbon dioxide
 (C) Both from water and carbon dioxide
 (D) Oxygen in air
- Q.3** First stable compound in C₃ cycle is
 (A) Phosphoglyceraldehyde
 (B) Phosphoglyceric acid
 (C) Fructose-1-6 diphosphate
 (D) Glucose-6-phosphate
- Q.4** Dark reaction of photosynthesis occurs in the
 (A) Stroma of the chloroplast outside the lamellae
 (B) Space between the two membranes of the chloroplast
 (C) Membranes of the stroma lamellae
 (D) Thylakoid membrane of the grana
- Q.5** A specific function of light energy in the process of photosynthesis is to
 (A) Activate chlorophyll
 (B) Oxidation of CO₂
 (C) Synthesis of glucose
 (D) Reduce CO₂
- Q.6** Digestion within a digestive tract is
 (A) Incomplete
 (B) Extracellular
 (C) The same as absorption
 (D) A reversible process
- Q.7** Dark reaction in photosynthesis is called so because
 (A) It does not require light energy
 (B) Cannot occur during daytime
 (C) Occurs more rapidly at night
 (D) It can also occur in darkness
- Q.8** Phloem always flows from a
 (A) Solar source to sugar sink
 (B) Sugar sink to sugar source
 (C) Leaf to the xylem to the phloem
 (D) Leaf to a root
- Q.9** With regards to natural eating habits, a human is
 (A) An herbivore (B) A carnivore
 (C) An omnivore (D) A Granivore
- Q.10** Muscular contractions of alimentary canal are
 (A) Circulation (B) Deglutition
 (C) Peristalsis (D) Churning
- Q.11** Which of the following regions of the alimentary canal of man does not secrete a digestive enzyme ?
 (A) Oesophagus (B) Stomach
 (C) Duodenum (D) Mouth
- Q.12** A digestive enzyme, salivary amylase, in the saliva begin digestion of
 (A) Protein (B) Nucleic acids
 (C) Fats (D) Carbohydrates
- Q.13** If you chew on a piece of bread long enough, it will begin to taste sweet because
 (A) Maltase is breaking down maltose
 (B) Lipases are forming fatty acids
 (C) Amylase is breaking down starches to disaccharides
 (D) Disaccharides are forming glucose
- Q.14** In the presence of lactase, lactose breaks down into molecules of
 (A) Glucose and galactose
 (B) Glucose and fructose
 (C) Galactose only
 (D) Glucose only
- Q.15** Saliva has the enzyme
 (A) Pepsin (B) Ptyalin
 (C) Trypsin (D) Rennin

- Q.16** Pepsin digests
 (A) Proteins in stomach
 (B) Carbohydrates in duodenum
 (C) Proteins in duodenum
 (D) Fats in ileum
- Q.17** Curding of milk in the stomach is due to the action of
 (A) Lipase (B) Rennin
 (C) Ptyalin (D) Tannin
- Q.18** Chief function of HCl is
 (A) To maintain a low pH to prevent growth of micro-organisms
 (B) To facilitate absorption
 (C) To maintain low pH to activate pepsinogen to form pepsin
 (D) To dissolve enzyme secreted in stomach
- Q.19** If the stomach did not producing any hydrochloric acid, which enzyme will not function ?
 (A) Ptyalin
 (B) Trypsin
 (C) Pepsin
 (D) Collagenase
- Q.20** Chief function of bile is
 (A) To digest fat by enzymatic action
 (B) To emulsify fat for digestion
 (C) To eliminate waste product
 (D) To regulate process of digestion
- Q.21** Where is bile produced ?
 (A) In gall bladder
 (B) In blood
 (C) In liver
 (D) In spleen
- Q.22** Ileum is
 (A) First part of the small intestine
 (B) Middle part of the small intestine
 (C) Last part of the small intestine
 (D) Not a part of the small intestine
- Q.23** Largest gland in human body is
 (A) Liver
 (B) Pancreas
 (C) Pituitary
 (D) Thyroid
- Q.24** The specific function of liver is
 (A) Excretion
 (B) Digestion
 (C) Histolysis
 (D) Glycogenesis and glycogenolysis
- Q.25** The original function of the vertebrate stomach was
 (A) Storage
 (B) Digestion
 (C) Enzyme secretion
 (D) Absorption
- Q.26** What is bile ?
 (A) a type of cell
 (B) a type of tooth
 (C) a solution that helps to break down fats
 (D) a food droplet
- Q.27** What do proteins get broken down into ?
 (A) amino acids (B) glucose
 (C) starch (D) fatty acids
- Q.28** What are the conditions in the stomach ?
 (A) alkaline (B) neutral
 (C) cold (D) acidic
- Q.29** What do large insoluble molecules get broken down into ?
 (A) large soluble molecules
 (B) small soluble molecules
 (C) medium soluble molecules
 (D) small insoluble molecules
- Q.30** When does photosynthesis take place ?
 (A) during the day (B) at night
 (C) night and day (D) in the winter

EXERCISE-3

- Q.1** Bile Juice is secreted from -
(Raj./NTSE Stage-1/Nutrition/13)
(A) Salivary glands
(B) Intestinal glands
(C) Stomach
(D) Liver
- Q.2** When acidity in Stomach increases, the medicine generally used is -
(Raj./NTSE Stage-1/Nutrition/13)
(A) Sodium bicarbonate
(B) Sodium Carbonate
(C) Ammonium Carbonate
(D) Ammonium bicarbonate
- Q.3** Substances necessary for autotrophic Nutrition are -
(Raj./NTSE Stage-1/Nutrition/13)
(A) CO₂ and H₂O (B) Chlorophyll
(C) Sun light (D) All of the above
- Q.4** Consider following chemical reaction
 $6\text{CO}_2 + 2\text{H}_2\text{O} \longrightarrow \text{C}_6\text{H}_{12}\text{O}_6 + 6\text{O}_2 + 6\text{H}_2\text{O}$
(Haryana/NTSE Stage-1/Nutrition/2013)
(a) CO₂ is oxidized to glucose
(b) CO₂ is reduced to glucose
(c) H₂O is oxidized to O₂
(d) H₂O is reduced to O₂
Which answer is correct?
(A) a & d (B) b & c
(C) b & d (D) a & c
- Q.5** Covering of lungs is -
(A) Pleura (B) Pericardium
(C) Epiglottis (D) Capsule
- Q.6** Exchange of gasses occurs through
(A) Stomata
(B) Lenticles
(C) Root surfaces
(D) All of the above
- Q.7** Concentrated nitric acid is used in a test to detect adulteration of :
(Karnataka /NTSE Stage-1/Nutrition/2013)
(A) Cooking oil (B) Milk
(C) Salt (D) Tea powder
- Q.8** In an experiment where two potted plants kept in a dark room are used to demonstrate that carbondioxide is essential for photosynthesis potassium hydroxide is used because it :
(Karnataka /NTSE Stage-1/Nutrition/2013)
(A) Releases oxygen
(B) Absorbs carbondioxide
(C) Releases carbon dioxide
(D) Absorbs oxygen
- Q.9** Which one of the following enzymes is present in saliva ?
(Mizoram/NTSE Stage-1/Nutrition/2013)
(A) Pepsin (B) Chymotrypsin
(C) Trypsin (D) Ptyalin
- Q.10** Which one of the following compound contains two carbon atoms ?
(Orissa/NTSE Stage-1/Nutrition/2013)
(A) Ethanol (B) Pyruvic acid
(C) Lactic acid (D) lucose
- Q.11** Read the following two statements and choose the correct answer.
(Orissa/NTSE Stage-1/Nutrition/2013)
i. Stomata regulates the body temperature in plants.
ii. Stomata helps in the absorption of minerals from the soil.
(A) i is true but ii is false
(B) i is false but ii is true
(C) Both i and ii are true
(D) Both i and ii are false
- Q.12** Which part of the human alimentary canal, is the site for complete digestion of carbohydrates.
(Punjab/NTSE Stage-1/Nutrition/2013)
(A) Stomach
(B) Small Intestine
(C) Large Intestine
(D) Rectum
- Q.13** This part of alimentary canal absorbs maximum amount of water and minerals:
(Maharashtra/NTSE Stage-1/Nutrition/2013)
(A) Small intestine
(B) Large intestine
(C) Stomach
(D) Oesophagus

- Q.14** Food becomes _____ due to bile juice.
(Maharashtra/NTSE Stage-1/Nutrition/2013)
(A) Acidic
(B) Alkaline
(C) Neutral
(D) First neutral and then acidic
- Q.15** Which of the following enzymes is related with digestion of protein ?
(Raj/NTSE Stage-1/Nutrition/2013)
(A) Lipase (B) Pepsin
(C) Sucrase (D) Amylase
- Q.16** What does liver secrete ?
(Gujarat/NTSE Stage-1/Nutrition/2013)
(A) Insulin (B) Bile
(C) Gastric juice (D) Mucus
- Q.17** Which of the following plant is a parasite ?
(Gujarat/NTSE Stage-1/Nutrition/2013)
(A) Cuscuta (B) Mushroom
(C) Giloe (D) Fern
- Q.18** Which two plant species obtain nutrition as symbionts in Lichens ?
(M.P./NTSE Stage-1/Nutrition/2013)
(A) Rhizobium and Drosera
(B) Fungi and Rose plant
(C) Algae and Virus
(D) Algae and Fungi
- Q.19** Assertion : Photosynthesis is minimum in green light
(Hariyanan/NTSE Stage-1/Nutrition/2013)
Reason : Chlorophylls are green in colour.
Direction :
(A) Both assertion and reason are true and reason is correct explanation of assertion.
(B) Both assertion and reason are true but reason is not correct explanation of assertion
(C) Assertion is true but reason is False
(D) Assertion is false but reason is true.
- Q.20** The substance essential for photosynthesis is
(Raj./NTSE Stage-1/Nutrition/2015)
(A) glucose (B) oxygen
(C) nitrogen (D) water
- Q.21** Chlorophyll contains.....
(M.P./NTSE Stage-1/Nutrition/2015)
(A) Potassium (B) Iron
(C) Manganese (D) Magnesium
- Q.22** Which of the following is an example of Insectivorous plant -
(M.P./NTSE Stage-1/Nutrition/2015)
(A) Cuscuta (B) Rafflesia
(C) Drosera (D) Tulsi
- Q.23** Photosynthesis is an important mode of autotrophic nutrition. The event which does not occur in photosynthesis is :
(Delhi/NTSE Stage-1/Nutrition/2015)
(A) Conversion of light energy to chemical energy
(B) Reduction of carbon dioxide to carbohydrate
(C) Oxidation of carbon to carbondioxide
(D) Absorption of light energy by chlorophyll
- Q.24** Which is the longest organ of the digestive system ?
(Gujarat/NTSE Stage-1/Nutrition/2015-16)
(A) Oesophagus (B) Stomach
(C) Small Intestine (D) Large Intestine
- Q.25** Deficiency of vitamin 'A' causes –
(M.P./NTSE Stage-1/Nutrition/2015-16)
(A) Beri-Beri (B) Anaemia
(C) Night blindness (D) Scurvy
- Q.26** One of the following juices secreted in the body of man does not contain any enzyme –
(Bihar/NTSE Stage-1/Nutrition/2015)
(A) Gastric juice (B) Saliva
(C) Bile juice (D) Pancreatic juice
- Q.27** Which of the following factors does a plant use for the process of photosynthesis ?
(Gujarat/NTSE Stage-1/Nutrition/2015-16)
(A) Sunlight (B) Chlorophyll
(C) CO₂ and H₂O (D) All of them
- Q.28** The substance not essential for photosynthesis is
(Raj./NTSE Stage-1/Nutrition/2017-18)
(A) sunlight (B) chlorophyll
(C) nitrogen (D) carbon dioxide.
- Q.29** The small intestine receives the secretion from _____
(Gujarat/NTSE Stage-1/2018-19)
(A) Salivary glands
(B) Stomach and liver
(C) Liver and salivary glands
(D) Liver and Pancreas

ANSWER KEY

EXERCISE - 2

Ques.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Ans.	B	B	B	A	A	B	A	A	C	C	A	D	C	A	B
Ques.	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
Ans.	A	B	C	C	B	C	C	A	D	A	C	A	D	B	A


EXERCISE - 3

Ques.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Ans.	D	A	D	B	A	D	B	B	D	A	A	B	B	B	B
Ques.	16	17	18	19	20	21	22	23	24	25	26	27	28	29	
Ans.	B	A	D	A	D	D	C	C	C	C	C	D	A	D	



REAL NUMBERS

Chapter Outline

- ✧ Number System
- ✧ Euclid's Division Lemma
- ✧ Euclid's Division Algorithm
- ✧ The Fundamental Theorem of Arithmetic
- ✧ Number of Factors
- ✧ HCF and LCM
- ✧ Logarithm
- ✧ Highest Power dividing Factorial




The Real Number Line



MIND MAP

REAL NUMBERS



NUMBER SYSTEM

- (i) A rational number is a number which can be put in the form $\frac{p}{q}$, where p and q are both integers and $q \neq 0$.
- (ii) A rational number is either a terminating or non-terminating and recurring (repeating) decimal.
- (iii) A rational number may be positive, negative or zero.

- (i) A number is irrational if and only if its decimal representation is non-terminating and non-repeating. e.g. $\sqrt{2}$, $\sqrt{3}$, π etc.
- (ii) Rational numbers and irrational number taken together form the set of real numbers.
- (iii) If a and b are two real numbers, then either (i) $a > b$ or (ii) $a = b$ or (iii) $a < b$
- (iv) Negative of an irrational number is an irrational number.
- (v) The sum of a rational number with an irrational number is always irrational.
- (vi) The product of a non-zero rational number with an irrational number is always an irrational number.
- (vii) The sum of two irrational numbers is not always an irrational number.
- (viii) The product of two irrational numbers is not always an irrational number.

Type (1) : Eg : $\frac{7}{8} = 0.875$

$$\begin{array}{r} 8 \overline{)70} \text{ (0.875)} \\ \underline{64} \\ 60 \\ \underline{56} \\ 40 \\ \underline{40} \\ \times \end{array}$$

∴ If decimal expansion ends (terminates) after finite number of steps, then decimal expansion of such numbers is terminating.

Eg : $\frac{1}{3} = 0.333.....$

$$= 0.\overline{3} \qquad \begin{array}{r} 3 \overline{)10} \overline{)0.33.....} \\ \underline{9} \\ 10 \\ \underline{9} \\ 1..... \end{array}$$

$$\text{or } \frac{1}{7} = 0.142857142857..... = 0.\overline{142857}$$

$$\begin{array}{r} 7 \overline{)10} \text{ (0.14285....} \\ \underline{7} \\ 30 \\ \underline{28} \\ 20 \\ \underline{14} \\ 60 \\ \underline{56} \\ 40 \\ \underline{35} \\ 50 \\ \underline{49} \\ 1.... \end{array}$$

In both examples remainder never becomes zero so the decimal expansion never ends after some or infinite steps of division. These type of decimal expansions are called non terminating.

Also we can see that on further division a digit of a block of digit repeats itself in the decimal part.

So these are called non terminating recurring decimal expansions.

Both the above types (1 & 2) are rational numbers.

Types (3) :

Eg :The decimal expansion 0.327172398... does not end any where, also there is no arrangement of digits (not repeating) so these are called non terminating not recurring.

These numbers are called irrational numbers.

Eg. :

0.1279312793	rational	terminating
0.1279312793....	rational	non terminating
or $0.\overline{12793}$		& recurring
0.32777	rational	terminating
$0.3\overline{27}$ or 0.32777.....	rational	non terminating & recurring
0.5361279	rational	terminating
0.10100100010000	rational	terminating
0.10100100010000...	irrational	non terminating non recurring.

Theorem-1 : Let p be a prime number. If p divides a^2 , then p divides a, where a is a positive integer.

Proof : Let the prime factorisation of a be as follows :

$a = p_1 p_2 p_n$, where p_1, p_2, p_n are primes, not necessarily distinct.

Therefore,

$$a^2 = (p_1 p_2 p_n) (p_1 p_2 p_n) = p_1^2 p_2^2 p_n^2 .$$

Now, we are given that p divides a^2 . Therefore, from the Fundamental Theorem of Arithmetic, it follows that p is one of the prime factors of a^2 . However, using the uniqueness part of the Fundamental Theorem of Arithmetic, we realise that the only prime factors of a^2 are $p_1, p_2, ..., p_n$. So p is one of $p_1, p_2, ..., p_n$.

Now, since $a = p_1 p_2 p_n$, p divides a.

Ex.1 Prove that $\sqrt{2}$ is irrational number

Sol. Let us assume, to contrary, that $\sqrt{2}$ is rational. That is, we can find integers a and b ($a, b \neq 0$) such that $\sqrt{2} = \frac{a}{b}$. Where a and b are coprime i.e. their H.C.F. = 1

So, $b\sqrt{2} = a$.

Squaring on both sides, and rearranging, we get $2b^2 = a^2$.

Therefore, a^2 is divisible by 2, and by Theorem 1, it follows that a is also divisible by 2.

So, we can write $a = 2c$ for some integer c .

Substituting for a , we get $2b^2 = 4c^2$,

That is, $b^2 = 2c^2$.

This means that b^2 is divisible by 2, and so b is also divisible by 2.

Therefore, a and b have at least 2 as a common factor.

But this contradicts the fact that a and b are coprime.

This contradiction has arisen because of our incorrect assumption that $\sqrt{2}$ is rational.

So, we conclude that $\sqrt{2}$ is irrational.

Euclid's Division Lemma

For any two positive integers a and b , there exist unique integers q and r satisfying $a = bq + r$, where $0 \leq r < b$.

For Example

(i) Consider number 23 and 5, then:

$$23 = 5 \times 4 + 3$$

Comparing with $a = bq + r$; we get:

$$a = 23, b = 5, q = 4, r = 3$$

and $0 \leq r < b$ (as $0 \leq 3 < 5$).

(ii) Consider positive integers 18 and 4.

$$18 = 4 \times 4 + 2$$

$\Rightarrow$ For $a = 18$ and $b = 4$ we have $q = 4$,

$$r = 2 \text{ and } 0 \leq r < b.$$

In the relation $a = bq + r$, where $0 \leq r < b$ is nothing but a statement of the long division of number a by number b in which q is the quotient obtained and r is the remainder.

Thus,

dividend = divisor $\times$ quotient + remainder

$$\Rightarrow a = bq + r$$

◆ H.C.F. (Highest Common Factor)

The H.C.F. of two or more positive integers is the largest positive integer that divides each given positive number completely.

For Example

(i) 14 is the largest positive integer that divides 28 and 70 completely; therefore H.C.F. of 28 and 70 is 14.

(ii) H.C.F. of 75, 125 and 200 is 25 as 25 divides each of 75, 125 and 200 completely and so on.

Euclid's Division Algorithm

If 'a' and 'b' are positive integers such that $a = bq + r$, then every common divisor of 'a' and 'b', is a common divisor of 'b' and 'r', and vice-versa.

- Using Euclid's Division Algorithm For Finding H.C.F.

Consider positive integers 418 and 33.

Step-1

Taking bigger number (418) as a and smaller number (33) as b

express the numbers as $a = bq + r$

$$\Rightarrow 418 = 33 \times 12 + 22$$

Step-2

Now taking the divisor 33 as a and remainder 22 as b apply the Euclid's division algorithm to get:

$$33 = 22 \times 1 + 11 \quad [\text{Expressing as } a = bq + r]$$

Step-3

Again with new divisor 22 as a and new remainder 11 as b apply the Euclid's division algorithm to get:

$$22 = 11 \times 2 + 0$$

Step-4

Since, the remainder = 0 so we cannot proceed further.

Step-5

The last divisor is 11 and we say H.C.F. of 418 and 33 is 11

Verification :

- (i) Using factor method:

$\therefore$ Factors of 418 = 1, 2, 11, 19, 22, 38, 209 and 418 and,

Factor of 33 = 1, 3, 11 and 33.

Common factors = 1 and 11

$$\Rightarrow \text{Highest common factor} = 11 \text{ i.e., H.C.F.} = 11$$

- (ii) Using prime factor method:

Prime factors of 418 = 2, 11 and 19.

Prime factors of 33 = 3 and 11.

$\therefore$ H.C.F. = Product of all common prime factors = 11. For any two positive integers a and b which can be expressed as, $a = bq + r$, where $0 \leq r < b$, the, H.C.F. of (a, b) = H.C.F. of (q, r) and so on. For number 418 and 33

$$418 = 33 \times 12 + 22$$

$$33 = 22 \times 1 + 11$$

and $22 = 11 \times 2 + 0$

$$\Rightarrow \text{H.C.F. of } (418, 33) = \text{H.C.F. of } (33, 22) = \text{H.C.F. of } (22, 11) = 11.$$

Ex.2 Using Euclid's division algorithm, find the H.C.F. of

(i) 135 and 225

(ii) 196 and 38220

- Sol.** (i) Starting with the larger number i.e., 225, we get : $225 = 135 \times 1 + 90$
 Now taking divisor 135 and remainder 90, we get : $135 = 90 \times 1 + 45$
 Further taking divisor 90 and remainder 45, we get : $90 = 45 \times 2 + 0$
 $\therefore$ Required H.C.F. = 45
- (ii) Starting with larger number 38220, we get : $38220 = 196 \times 195 + 0$
 Since, the remainder is 0
 $\Rightarrow$ H.C.F. = 196

Ex.3 Show that any positive odd integer is of the form $4q + 1$ or $4q + 3$, where q is some integer.

Sol. Let a and b be two positive integers where a is greater than b . According to Euclid's division lemma; a and b can be expressed as

$$a = bq + r, \text{ where } q \text{ is quotient and } r \text{ is remainder and } 0 \leq r < b.$$

Taking $b = 4$, we get

$$a = 4q + r,$$

where $0 \leq r < 4$ i.e., $r = 0, 1, 2$ or 3

$r = 0 \Rightarrow a = 4q$, which is divisible by 2 and so is even.

$r = 1 \Rightarrow a = 4q + 1$, which is not divisible by 2 and so is odd.

$r = 2 \Rightarrow a = 4q + 2$, which is divisible by 2 and so is even.

$r = 3 \Rightarrow a = 4q + 3$, which is not divisible by 2 and so is odd.

$\therefore$ Any positive odd integer is of the form

$4q + 1$ or $4q + 3$; where q is an integer.

Ex.4 Use Euclid's Division Algorithm to show that the square of any positive integer is either of the form $3m$ or $3m + 1$ for some integer m .

Sol. Let a and b are two positive integers such that a is greater than b ; then:

$$a = bq + r; \text{ where } q \text{ and } r \text{ are also positive integers and } 0 \leq r < b$$

Taking $b = 3$, we get:

$$a = 3q + r; \text{ where } 0 \leq r < 3$$

$\Rightarrow$ The value of positive integer a will be $3q + 0$, $3q + 1$ or $3q + 2$

i.e., $3q$, $3q + 1$ or $3q + 2$.

Now we have to show that the squares of positive integers $3q$, $3q + 1$ and $3q + 2$ can be expressed as $3m$, or $3m + 1$ for some integer m .

$$\therefore \text{ Square of } 3q = (3q)^2$$

$$= 9q^2 = 3(3q^2) = 3m; \text{ where } m \text{ is some integer.}$$

$$\text{Square of } 3q + 1 = (3q + 1)^2$$

$$= 9q^2 + 6q + 1$$

$$= 3(3q^2 + 2q) + 1 = 3m + 1 \text{ for some integer } m.$$

$$\text{Square of } 3q + 2 = (3q + 2)^2$$

$$= 9q^2 + 12q + 4$$

$$= 9q^2 + 12q + 3 + 1$$

$$= 3(3q^2 + 4q + 1) + 1 = 3m + 1 \text{ for some integer } m.$$

$\therefore$ The square of any positive integer is either of the form $3m$ or $3m + 1$ for some integer m .

COMPETITIVE LEVEL

Note :

- (i) $(x^n - a^n)$ is divisible by $(x - a)$ for all the values of n .
- (ii) $(x^n - a^n)$ is divisible by $(x + a)$ for all even values of n .
- (iii) $(x^n + a^n)$ is divisible by $(x + a)$ for all odd values of n .

◆ Euclid Division Algorithm

$$\text{Dividend} = (\text{Divisor} \times \text{Quotient}) + \text{Remainder}$$

Ex.5 On dividing 20032 by a certain number, the quotient is 101 and the remainder is 135. Find the divisor.

Sol. $\text{Divisor} = \frac{\text{Dividend} - \text{Remainder}}{\text{Quotient}} = \frac{20032 - 135}{101} = 197.$

Test of Divisibility :

No.	Divisibility Test
2	Unit digit should be 0 or even.
3	The sum of digits of no. should be divisible by 3.
4	The no. formed by last 2 digits of given no. should be divisible by 4.
5	Unit digit should be 0 or 5.
6	No. should be divisible by 2 & 3 both.
7	No. without ones - 2(ones) = no. should be 0 or divisible by 7.
8	The number formed by last 3 digits of given no. should be divisible by 8.
9	Sum of digits of given no. should be divisible by 9.
11	The difference between sums of the digits at even & at odd places should be zero or multiple of 11.
13	No. without ones + 4 (ones digit) = no. should be divisible by 13.
17	No. without ones - 5(ones) = no. should be 0 or divisible by 17.
19	No. without ones + 2(ones) = no. should be divisible by 19.
25	Last 2 digit of the number should be 00, 25, 50 or 75.

The Fundamental Theorem Of Arithmetic

Statement : Every composite number can be decomposed as product of prime numbers in a unique way, except for the order in which the prime numbers occur.

For example :

(i) $30 = 2 \times 3 \times 5, 30 = 3 \times 2 \times 5, 30 = 2 \times 5 \times 3$ and so on.

(ii) $432 = 2 \times 2 \times 2 \times 2 \times 3 \times 3 \times 3 = 2^4 \times 3^3$

or $432 = 3^3 \times 2^4.$

(iii) $12600 = 2 \times 2 \times 2 \times 3 \times 3 \times 5 \times 5 \times 7 = 2^3 \times 3^2 \times 5^2 \times 7$

In general, a composite number is expressed as the product of its prime factors written in ascending order of their values.

E.g., (i) $6615 = 3 \times 3 \times 3 \times 5 \times 7 \times 7 = 3^3 \times 5 \times 7^2$

(ii) $532400 = 2 \times 2 \times 2 \times 2 \times 5 \times 5 \times 11 \times 11 \times 11 = 2^4 \times 5^2 \times 11^3$

COMPETITIVE LEVEL**HCF and LCM of fractions :**

$$\text{LCM of fractions} = \frac{\text{LCM of numerators}}{\text{HCF of denominators}}$$

$$\text{HCF of fractions} = \frac{\text{HCF of numerators}}{\text{LCM of denominators}}$$

Make sure the fractions are in the most reducible form.

Ex.15 Find a number which when divided by 3, 4, and 13 always leaves the same remainder 2.

Sol. The smallest number which, when divided by 3, 4 and 13, leaves the remainder 2 in each case is $\text{LCM}(3, 4 \text{ and } 13) + 2 = 156 + 2 = 158$

Ex.16 Find the H.C.F. and L.C.M. of $\frac{2}{3}$, $\frac{8}{9}$, $\frac{16}{81}$ and $\frac{10}{27}$.

$$\text{Sol. H.C.F. of given fractions} = \frac{\text{H.C.F. of } (2, 8, 16, 10)}{\text{L.C.M. of } (3, 9, 81, 27)} = \frac{2}{81},$$

$$\text{L.C.M. of given fractions} = \frac{\text{L.C.M. of } (2, 8, 16, 10)}{\text{H.C.F. of } (3, 9, 81, 27)} = \frac{80}{3}.$$

Logarithm

If 'a' is a positive real number, other than 1 and x is a real number such that $a^x = N$, then x is the logarithm of N to the base a.

$$\text{If } a^x = N \text{ then } \log_a N = x.$$

Where $N > 0$ and $a > 0, a \neq 1$

Systems of Logarithm

There are two systems of logarithm which are generally used.

- (i) Common logarithm : In this system base is always taken as 10.
- (ii) Natural logarithm : In this system the base of the logarithm is taken as 'e'. Where 'e' is an irrational number lying between 2 and 3. (The approximate value of e upto two decimal places is equal to 2.73)

Fundamental Laws of Logarithm

Logarithm to any base a (where $a > 0$ and $a \neq 1$).

$$(i) \log_a a = 1$$

$$(ii) \log_a 0 = \text{not defined} \quad [\text{As } a^n = 0 \text{ is not possible, where } n \text{ is any number}]$$

$$(iii) \log_a (-\text{ve no.}) = \text{not defined.} \quad [\text{As in } \log_a N, N \text{ will always be (+ve)}]$$

$$(iv) \log_a (mn) = \log_a m + \log_a n \quad [\text{Where } m \text{ and } n \text{ are +ve numbers}]$$

$$(v) \log_a \left(\frac{m}{n} \right) = \log_a m - \log_a n$$

$$(vi) \log_a (m)^n = n \log_a m$$

$$(vii) \log_a m = \frac{\log_b m}{\log_b a}$$

$$(viii) \log_a m \cdot \log_m a = 1$$

$$(ix) a^{\log_a n} = n$$

$$(x) \log_{a^q} n^p = \frac{p}{q} \log_a n, n > 0$$

$$(xi) p^{\log_a q} = q^{\log_a p}, n > 0$$

$$(xii) \log_a x = \log_a y, x = y$$

Ex.17 Find the value of $\log \frac{9}{8} - \log \frac{27}{32} + \log \frac{3}{4}$

Sol. Given :

$$\begin{aligned} \log \frac{9}{8} - \log \frac{27}{32} + \log \frac{3}{4} &= \log \left(\frac{9}{8} \div \frac{27}{32} \right) + \log \frac{3}{4} \\ &= \log \left(\frac{9}{8} \times \frac{32}{27} \times \frac{3}{4} \right) = \log 1 = 0. \quad [\log_a 1 = 0] \end{aligned}$$

Ex.18 If $2\log_4 x = 1 + \log_4(x - 1)$, find the value of x .

Sol. Given $2\log_4 x = 1 + \log_4(x - 1)$

$$\Rightarrow \log_4 x^2 - \log_4(x - 1) = 1$$

$$\Rightarrow \log_4 \frac{x^2}{x - 1} = 1$$

$$\Rightarrow 4^1 = \frac{x^2}{x - 1}$$

$$\Rightarrow x^2 = 4x - 4$$

$$\Rightarrow x^2 - 4x + 4 = 0$$

$$\Rightarrow (x - 2)^2 = 0 \quad x = 2.$$

Ex.19 If $A = \log_{27} 625 + 7^{\log_{11} 13}$ and $B = \log_9 125 + 13^{\log_{11} 7}$, then find the relation between A and B .

Sol. $A = \log_{27} 625 + 7^{\log_{11} 13} = \log_{3^3} 5^4 + 7^{\log_{11} 13}$

$$\text{or, } A = \frac{4}{3} \log_3 5 + 7^{\log_{11} 13} \quad \dots(i)$$

$$\text{and, } B = \log_9 125 + 13^{\log_{11} 7}$$

$$\text{or, } B = \log_{3^2} 5^3 + 7^{\log_{11} 13}$$

$$\text{or, } B = \frac{3}{2} \log_3 5 + 7^{\log_{11} 13} \quad \dots(ii)$$

By (i) and (ii) we have,

$$A - \frac{4}{3} \log_3 5 = B - \frac{3}{2} \log_3 5$$

$$\therefore \frac{4}{3} \log_3 5 < \frac{3}{2} \log_3 5 \quad \therefore A < B.$$

Highest Power Dividing Factorial

Factorial n : Product of n consecutive natural numbers starting from 1 is known as 'factorial n ' it is denoted by ' $n!$ '.

So, $n! = n.(n-1).(n-2) \dots 3.2.1$. e.g. $5! = 5 \times 4 \times 3 \times 2 \times 1 = 120$.

The value of factorial zero is equal to the value of factorial one. Hence $0! = 1 = 1!$

The approach to finding the highest power of x dividing $y!$ is $\left[\frac{y}{x} \right] + \left[\frac{y}{x^2} \right] + \left[\frac{y}{x^3} \right] \dots$, where $[]$ represents just the integral part of the answer and ignoring the fractional part.

Ex.20 What is the highest power of 2 that divides $20!$ completely?

Sol. $20! = 1 \times 2 \times 3 \times 4 \times \dots \times 18 \times 19 \times 20 = 1 \times (2^1) \times 3 \times (2^2) \times 5 \times (2^1 \times 3^1) \times 7 \times (2^3) \times \dots$ so on. In order to find the highest power of 2 that divides the above product, we need to find the sum of the powers of all 2 in this expansion. All numbers that are divisible by 2^1 will contribute 1 to the exponent of 2 in the product = 10. Hence, 10 numbers contribute 2^1 to the product. Similarly, all numbers that are divisible by 2^2 will contribute an extra 1 to the exponent of 2 in the product, i.e. $\frac{20}{2^2} = 5$. Hence, 5 numbers contribute an extra 1 to exponents. Similarly, there are 2 numbers that are divisible by 2^3 and 1 number that is divisible by 2^4 . Hence, the total 1s contributed to the exponent of 2 in $20!$ is the sum of $(10 + 5 + 2 + 1) = 18$. Hence, group of all 2s in $20!$ gives $2^{18} \times (N)$, where N is not divisible by 2.

If $20!$ is divided by 2^x then maximum value of x is 18.

Ex.21 How many zeros at the end of first 100 multiples of 10.

Sol. First 100 multiple of 10 are $= 10 \times 20 \times 30 \times \dots \times 1000$

$$= 10^{100} (1 \times 2 \times 3 \times \dots \times 100)$$

$$= 10^{100} \times 10^{24} \times N$$

$$= 10^{124} \times N$$

Where N is not divisible by 10

So, there are 124 zero at the end of first 100 multiple of 10.

EXERCISE-1

➤ Very Short Answer Type Questions

Q.1 Classify the following numbers as rational or irrational :

- | | |
|-----------------------|----------------------------|
| (i) $\frac{22}{7}$ | (ii) 3.1416 |
| (iii) π | (iv) $\overline{3.142857}$ |
| (v) 5.636363..... | (vi) 2.040040004..... |
| (vii) 1.535335333.... | (viii) 3.121221222... |
| (ix) $\sqrt{21}$ | (x) $\sqrt[3]{3}$ |

Q.2 Prove that $\frac{1}{\sqrt{3}}$ is irrational.

Q.3 Define Euclid's Division Lemma.

Q.4 A number when divided by 61 gives 27 as quotient and 32 as remainder. Find the number.

Q.5 By what number should 1365 be divided to get 31 as quotient and 32 as remainder ?

➤ Short Answer Type Questions – Type I

Q.6 Find two irrational numbers between 3 and 4.

Q.7 Find two irrational numbers between 2 and 2.5.

Q.8 Consider the number 6^n , where n is a natural number. Check whether there is any value of $n \in \mathbb{N}$ for which 6^n is divisible by 7.

Q.9 Prove that each of the following numbers is irrational :

- | | |
|------------------------------|--------------------------|
| (i) $\sqrt{6}$ | (ii) $(2 - \sqrt{3})$ |
| (iii) $(3 + \sqrt{2})$ | (iv) $(2 + \sqrt{5})$ |
| (v) $(5 + 3\sqrt{2})$ | (vi) $3\sqrt{7}$ |
| (vii) $\frac{3}{\sqrt{5}}$ | (viii) $(2 - 3\sqrt{5})$ |
| (ix) $(\sqrt{3} + \sqrt{5})$ | |

Q.10 Without actual division, show that each of the following rational numbers is a non-terminating repeating decimal :

- | | |
|---------------------------------|--|
| (i) $\frac{11}{(2^3 \times 3)}$ | (ii) $\frac{73}{(2^3 \times 3^3 \times 5)}$ |
| (iii) $\frac{9}{35}$ | (iv) $\frac{32}{147}$ |
| (v) $\frac{64}{455}$ | (vi) $\frac{77}{210}$ |
| (vii) $\frac{29}{343}$ | (viii) $\frac{129}{(2^2 \times 5^7 \times 7^5)}$ |

➤ Short Answer Type Questions – Type II

Q.11 Without actual division, show that each of the following rational numbers is a terminating decimal. Express each in decimal form :

- | | | |
|-----------------------------------|-----------------------|------------------------|
| (i) $\frac{23}{(2^3 \times 5^2)}$ | (ii) $\frac{24}{125}$ | (iii) $\frac{17}{320}$ |
| (iv) $\frac{171}{800}$ | (v) $\frac{15}{1600}$ | (vi) $\frac{19}{3125}$ |

Q.12 Express each of the following as a fraction in simplest form :

- | | | |
|------------------------|-----------------------|-------------------------|
| (i) $0.\overline{8}$ | (ii) $2.\overline{4}$ | (iii) $0.\overline{24}$ |
| (iv) $0.\overline{12}$ | (v) $2.\overline{24}$ | (vi) $0.\overline{365}$ |

Q.13 Using prime factorisation, find the HCF and LCM of

- | | |
|------------------|---------------------|
| (i) 144, 198 | (ii) 396, 1080 |
| (iii) 1152, 1664 | (iv) 21, 28, 36, 45 |

Q.14 The HCF of two numbers is 23 and their LCM is 1449. If one of the numbers is 161, find the other.

Q.15 Find the $[\text{HCF} \times \text{LCM}]$ for the numbers 105 and 120.

Q.16 Three pieces of timber 42 m, 49 m and 63 m long have to be divided into planks of the same length. What is the greatest possible length of each plank ?

- Q.17** Find the greatest possible length which can be used to measure exactly the length 7 m, 3 m 85 cm and 12 m 95 cm.
- Q.18** Three sets of English, Mathematics and Science books containing 336, 240 and 96 books respectively have to be stacked in such a way that all the books are stored subject wise and the height of each stack is the same. How many stacks will be there ?
- Q.19** Three measuring rods are 64 cm, 80 cm and 96 cm in length. Find the least length of cloth that can be measured an exact number of times, using any of the rods.
- Q.20** The traffic lights at three different road crossings change after every 48 seconds, 72 seconds and 108 seconds respectively. If they all change simultaneously at 8 a.m., then at what time will they again change simultaneously ?

➤ Long Answer Type Questions

- Q.21** An electronic device makes a beep after every 60 seconds. Another device makes a beep after every 62 seconds. They beeped together at 10 am. At what time will they beep together at the earliest ?
- Q.22** Using Euclid's algorithm, find the HCF of
(i) 405 and 2520
(ii) 504 and 1188
(iii) 960 and 1575
- Q.23** Show that any positive integer which is of the form $6q + 1$ or $6q + 3$ or $6q + 5$ is odd, where q is some integer.
- Q.24** Show that one and only one out of n , $n + 2$ or $n + 4$ is divisible by 3, where n is any positive integer
- Q.25** Find the least number of square tiles required to pave the ceiling of a room 15 m 17 cm long and 9 m 2 cm broad.
- Q.26** Six bells commence tolling together and toll at intervals of 2, 4, 6, 8, 10, 12 minutes respectively. In 30 hours, how many times do they toll together ?
- Q.27** Show that there is no positive integer n , for which $\sqrt{n-1} + \sqrt{n+1}$ is rational.
- Q.28** Find HCF of 81 and 237 and express it as a linear combination of 81 and 237 i.e., $\text{HCF}(81, 237) = 81x + 237y$ for some x and y . Find the value of x and y .
- Q.29** Three friends Salman, Hritik & John were good friends. They need to go for morning walk together, on a morning walk, they step off together & their steps measure 40 cm, 42 cm, & 45 cm respectively.
(a) What is minimum distance each should walk so that each can cover same distance in complete steps ?
(b) What you have learnt (values/lesson) from above activity of three friends.
- Q.30** Aakriti decided to distribute milk in an orphanage on her birthday. The supplier brought two milk containers which contain 398 l and 436 l of milk. The milk is to be transferred to another containers so 7 l and 11 l of milk is left in both the containers respectively
(a) What will be the maximum capacity of the drum?
(b) What qualities/ values were shown by Aakriti ?

EXERCISE-2

- Q.1** The product of the digits of a four digit number is 540. If the digits are all different find the greatest digit out of them if the least is 2.
(A) 5 (B) 6
(C) 9 (D) None of these
- Q.2** How many natural numbers are there which satisfy $n^2 < 8n < 35n < n^3$.
(A) 1 (B) 2 (C) 3 (D) 4
- Q.3** xy is a number that is divided by ab where $xy < ab$ and gives a result $0.xyxyxy\dots$ then ab equals :
(A) 11 (B) 33 (C) 99 (D) 66
- Q.4** Given that a, b are odd and c, d are even, then
(A) $a^2 - b^2 + c^2 - d^2$ is always divisible by 4
(B) $abc + bcd + cda + dab$ is always divisible by 4
(C) $a^4 + b^4 + c^3 + d^3 + c^2b + a^2b$ is always odd
(D) $a + 2b + 3c + 4d$ is odd
- Q.5** If $a, a + 2$, and $a + 4$ are prime numbers, then the number of possible solution for a is -
(A) three (B) two
(C) one (D) more than three
- Q.6** There are four prime numbers written in ascending order. The product of the first three is 385 and that of the last three is 1001. The last number is :
(A) 11 (B) 13 (C) 17 (D) 19
- Q.7** In a six digit number 5 digits are prime numbers. The sum of all the digits is 24. The 2nd, 3rd and 5th digit are identical and the others are distinct digits. The number is divisible by 4. The last digit of the number is
(A) 2 or 4 (B) 4 or 6
(C) 4 or 6 or 8 (D) 2 or 6 or 8
- Q.8** The number of positive n in the range $12 \leq n \leq 40$ such that the product $(n - 1)(n - 2)\dots 3.2.1$ is not divisible by n is :
(A) 5 (B) 7 (C) 13 (D) 14
- Q.9** If least prime factor of a number m is 3 and least prime factor of another number n is 7, then least prime factor of the number $(m + n)$ is :
(A) 2 (B) 3 (C) 5 (D) 7
- Q.10** V is product of first 41 natural numbers. $A = V + 1$. The number of primes among $A + 1, A + 2, A + 3, A + 4, \dots, A + 39, A + 40$ is :
(A) 1 (B) 2 (C) 3 (D) 0
- Q.11** If N denotes the number of digits in the number $(5^{84})(2^{86})$ then N equals
(A) 83 (B) 84
(C) 85 (D) 88
- Q.12** How many pairs of positive integers (n, m) , with $n \leq m$ satisfy the equation $\frac{1}{5} = \frac{1}{n} + \frac{1}{m}$?
(A) 1 (B) 2 (C) 3 (D) 4
- Q.13** If $a^2 - b^2 = 13$ where a and b are natural numbers, then value of a is :
(A) 6 (B) 7 (C) 8 (D) 9
- Q.14** Find the number of divisors of $2^7 \times 3^8 \times 5^9$ -
(A) 504 (B) 720
(C) 724 (D) 540
- Q.15** Find the least number which when divided by 15, leaves a remainder of 5, when divided by 25, leaves a remainder of 15 and when divided by 35 leaves a remainder of 25.
(A) 515 (B) 525
(C) 1040 (D) 1050
- Q.16** If $\text{HCF}(p, q) = 12$ and $p \times q = 1800$ then $\text{LCM}(p, q)$ is :
(A) 3600 (B) 900
(C) 150 (D) 90

- Q.17** The least number of square tiles required to pave the ceiling of a room 15 m 17 cm long and 9 m 2 cm broad, is :
(A) 902 (B) 656 (C) 738 (D) 814
- Q.18** Four watches are ringing alarm bells in the interval of 6, 12, 15 and 18 seconds. If they start at same time, how many times they will ring together in 4 hrs.
(A) 80 times (B) 81 times
(C) 20 times (D) 21 times
- Q.19** If $4^{25} + 5^{15}$ is divided by 10, then the remainder is :
(A) 4 (B) 5 (C) 9 (D) 0
- Q.20** Number of values of 'a' (from 0 to 9) for the number $N = 2345631143a4$ is divisible by 12, is :
(A) 0 (B) 1
(C) 2 (D) more than 2
- Q.21** $1^{13} + 2^{13} + 3^{13} + \dots + 60^{13}$ is divisible by :
(A) 61 (B) 63
(B) 65 (D) 59
- Q.22** The value of the digit d for which the number d456d is divisible by 18, is :
(A) 3 (B) 4 (C) 6 (D) 9
- Q.23** The smallest integral value of x, for which $\frac{7}{x}$ is an integer is :
(A) 1 (B) -1 (C) 7 (D) -7
- Q.24** $19^n - 1$ is -
(A) always divisible by 9
(B) always divisible by 20
(C) is never divisible by 19
(D) only (A) and (C) are true
- Q.25** Which one of the numbers listed below is not a divisor of the number $N = (2^{30} - 1)$, is equal to :
(A) $2^5 - 1$ (B) $2^5 + 1$
(C) $2^6 - 1$ (D) $2^{10} + 1$
- Q.26** $8^n - 6^n - 3^n + 1$ is a multiple of 10 :
(A) 5 (B) 2
(C) 10 (D) All of these
- Q.27** The least number which on division by 35 leaves a remainder 25 and on division by 45 leaves the remainder 35 and on division by 55 leaves the remainder 45 is:
(A) 2515 (B) 3455
(C) 2875 (D) 2785
- Q.28** When $4^{101} + 6^{101}$ is divided by 25, the remainder is :
(A) 20 (B) 10 (C) 5 (D) 0
- Q.29** What is the remainder when 13^{400} is divided by 11?
(A) -1 (B) 1
(C) 5 (D) 2
- Q.30** The greatest value assumed by the function $f(x) = 5 - |x - 3|$ is :
(A) 3 (B) 8 (C) 6 (D) 5
- Q.31** If $x < -2$, then $|1 - |1 + x||$ equals :
(A) $2 + x$ (B) x
(C) $-x$ (D) $-(2 + x)$
- Q.32** If $\log_{10} 2 = 0.30103$, $\log_{10} 3 = 0.47712$, the number of digits in $3^{12} \times 2^8$ is
(A) 7 (B) 8 (C) 9 (D) 10
- Q.33** The value of $\log_3 4 \log_4 5 \log_5 6 \log_6 7 \log_7 8 \log_8 9$ is
(A) 1 (B) 2 (C) 3 (D) 4
- Q.34** The value of $81^{(1/\log_5 3)} + 27^{\log_9 36} + 3^{4/\log_7 9}$ is equal to -
(A) 49 (B) 625
(C) 216 (D) 890
- Q.35** If $\log_4 5 = a$ and $\log_5 6 = b$, then $\log_3 2$ is equal to
(A) $\frac{1}{2a+1}$ (B) $\frac{1}{2b+1}$
(C) $2ab + 1$ (D) $\frac{1}{2ab-1}$
- Q.36** $\log_{10} p + \log_{10} q = \log_{10} (p-q)$, then :
(A) $p = q = 0$ (B) $p = \frac{q}{1+q}$
(C) $p = q = 1$ (D) $p = \frac{q}{1-q}$

- Q.37** Find the number of positive integers which divide 10^{999} but not 10^{998} .
 (A) 1999 (B) 1
 (C) 999 (D) None of these

- Q.38** The power of 2 in 150!
 (A) 150 (B) 146
 (C) 140 (D) 156

- Q.39** Find all real numbers x satisfying
 $x = 2(2(2(2x - 1) - 1) - 1) - 1$
 (A) 15 (B) 2
 (C) 3 (D) None of these

- Q.40** $\frac{1}{2} + \frac{2^1}{2^2} + \frac{2^2}{2^3} + \dots + \frac{2^{2006}}{2^{2007}} + \frac{2^{2007}}{2^{2008}}$ is equal to :
 (A) 1004 (B) $\frac{1}{2^{2008}}$
 (C) 2008 (D) 502

- Q.41** If n is a natural number then we define $n!$ (pronounced as factorial n) to be the product $n \times (n - 1) \times (n - 2) \times \dots \times 2 \times 1$. For example $4! = 4 \times 3 \times 2 \times 1 = 24$. If $6! = a! \times b!$ where $a > 1$ and $b > 1$, then $a + b$ is :
 (A) 8 (B) 7 (C) 6 (D) 5

- Q.42** In the sequencea, b, c, d, 0, 1, 1, 2, 3, 5, 8.... each term is the sum of the two terms to its left. Find 'a'.
 (A) -3 (B) -1
 (C) 0 (D) 1

- Q.43** The value of $\left(1 - \frac{1}{3}\right)^2 \left(1 - \frac{1}{4}\right)^2 \left(1 - \frac{1}{5}\right)^2 \dots \left(1 - \frac{1}{n}\right)^2$ is equal to :
 (A) $\left(\frac{1}{n}\right)^2$ (B) $\left(\frac{2}{n}\right)^2$
 (C) $\left(\frac{3}{n}\right)^2$ (D) $\left(\frac{4}{n}\right)^2$

- Q.44** The value of the expression
 $\sqrt{\frac{1}{\sqrt{2}+1} + \frac{1}{\sqrt{3}+\sqrt{2}} + \frac{1}{\sqrt{4}+\sqrt{3}} + \dots \text{upto 99 terms}}$
 is equal to :
 (A) 9 (B) 3 (C) 1 (D) 0

- Q.45** The number $N = \sqrt[3]{2+\sqrt{5}} + \sqrt[3]{2-\sqrt{5}}$ equals :
 (A) 1 (B) $\sqrt{5} - 1$
 (C) $\sqrt[3]{2}$ (D) $\sqrt{5} - \sqrt[3]{2}$

- Q.46** Find the value of
 $^{507}\sqrt{2\sqrt{7}-3\sqrt{3}} \cdot ^{1014}\sqrt{55+12\sqrt{21}}$
 (A) 1 (B) 2
 (C) 5 (D) None of these

- Q.47** If $\sqrt{9-(n-2)^2}$ is a real number, then the number of integral values of n is :
 (A) 3 (B) 5
 (C) 7 (D) Infinitely many

- Q.48** If x is a positive integer less than 100, then the number of x which make $\sqrt{1+2+3+4+\dots+x}$ an integer is:
 (A) 6 (B) 7 (C) 8 (D) 9

- Q.49** If $A^{\frac{1}{A}} = B^{\frac{1}{B}} = C^{\frac{1}{C}}$, $A^{BC} + B^{AC} + C^{AB} = 729$. Which of the following equals $A^{\frac{1}{A}}$?
 (A) $^{ABC}\sqrt{81}$ (B) $^{ABC}\sqrt{243}$
 (C) $^{ABC}\sqrt{27}$ (D) $^{ABC}\sqrt{9}$

- Q.50** For $x \neq 0, \pm 1$, the expression
 $\frac{\frac{1}{x^{2007}} - \frac{1}{x^{2009}}}{\frac{1}{x^{2008}} - \frac{1}{x^{2010}}}$ is equivalent to :
 (A) x (B) $x - 1$
 (C) $x^2 - 1$ (D) $\frac{1}{x}$

EXERCISE-3

- Q.1** If a and b are natural numbers such that $\left(\frac{1}{a}\right)^b = 0.\bar{3}$, then the value of ab is :
[NTSE Stage-II 2007]
(A) 81 (B) 24 (C) 192 (D) 375
- Q.2** If the sum of the digits of a number $(10^n - 1)$ is 4707, where n is a natural number, then the value of n is :
[NTSE Stage-II/2007]
(A) 477 (B) 523 (C) 532 (D) 704
- Q.3** If $5\frac{7}{x} \times y\frac{1}{13} = 12$, where fractions are in their lowest terms, then $x - y$ is equal to
[NTSE Stage-II /2010]
(A) 2 (B) 4 (C) 7 (D) 9
- Q.4** $1x3y6$ is a five digit number where x, y are digits and y exceeds x by 6. If this number is divisible by 18, then the value of $\frac{y}{x}$ is :
[NTSE Stage-II/2010]
(A) 7 (B) 3 (C) $\frac{1}{3}$ (D) $\frac{1}{7}$
- Q.5** The sum of any three distinct natural numbers arranged in ascending order is 200 such that the second number is a perfect cube. How many possible values are there for this number ?
[NTSE Stage-II/2011]
(A) 4 (B) 3 (C) 2 (D) 1
- Q.6** If the digits of a three-digit number are reversed, then the number so obtained is less than the original number by 297. If the sum of the digit of the number is 8 and its hundred's digit has the largest possible value, then the ten's digit of the number is :
[NTSE Stage-II/2011]
(A) 3 (B) 2 (C) 1 (D) 0
- Q.7** Number of zero's in the product of $5 \times 10 \times 25 \times 40 \times 50 \times 55 \times 65 \times 125 \times 80$, is -
[M.P. NTSE Stage-I 2013]
(A) 8 (B) 9 (C) 12 (D) 13
- Q.8** A farmer divides his herd of x cows among his 4 son's such that first son gets one-half of the herd, the second son gets one fourth, the third son gets one-fifth and the fourth son gets 7 cows, then the value of x is :
[M.P. NTSE Stage-I 2013]
(A) 100 (B) 140 (C) 160 (D) 180
- Q.9** If $\log_{12} 27 = a$, then $\log_6 16$ is
[Delhi NTSE Stage-I 2013]
(A) $\frac{4(3-a)}{(3+a)}$ (B) $\frac{4(3+a)}{(3-a)}$
(C) $\frac{(3+a)}{4(3-a)}$ (D) $\frac{3-a}{4(3+a)}$
- Q.10** The value of $4\log\frac{24}{25} - 16\log\frac{9}{10} + 7\log\frac{81}{80}$
[M.P. NTSE Stage-I 2014]
(A) $\log 9$ (B) $\log 7$
(C) $\log 5$ (D) $\log 3$
- Q.11** Which real number lies between 2 and 2.5 : [Chandigarh NTSE Stage-I 2014]
(A) $\sqrt{11}$ (B) $\sqrt{8}$
(C) $\sqrt[3]{7}$ (D) $\sqrt[3]{9}$
- Q.12** For positive x and y , the LCM is 225 and HCF is 15 There. [NTSE Stage-II 2014]
(A) is exactly one such pair
(B) are exactly two such pair
(C) are exactly three such pair
(D) are exactly four such pair
- Q.13** The HCF of any two prime numbers a and b , is -
[Rajasthan NTSE Stage-I 2015]
(A) a (B) ab (C) b (D) 1

- Q.14** The traffic lights at three different signals change after 48 seconds, 72 seconds and 108. If they change at 7 a.m. simultaneously. How many times they will change between 7 a.m. to 7 : 30 a.m. simultaneously ?

[Haryana NTSE Stage-I 2015]

- (A) 3 (B) 4 (C) 5 (D) 2

- Q.15** The smallest number which when increased by 17 is exactly divisible by both 520 and 468 is :

[Karnataka NTSE stage - I 2014]

- (A) 4697 (B) 4656
(C) 4663 (D) 4680

- Q.16** Which of the following is an irrational number? [NTSE Stage-II 2013]

- (A) $\sqrt{41616}$
(B) 23.232323 ...
(C) $\frac{(1+\sqrt{3})^3 - (1-\sqrt{3})^3}{\sqrt{3}}$
(D) 23.10100100010000 ...

- Q.17** On dividing a natural number by 13, the remainder is 3 and on dividing the same number by 21, the remainder is 11. If the number lies between 500 and 600, then the remainder on dividing the number by 19 is - [NTSE Stage-II 2016]

- (A) 4 (B) 6 (C) 9 (D) 13

- Q.18** Expressing $0.\overline{34} + 0.\overline{34}$ as a single decimal, we get - [NTSE Stage-II 2016]

- (A) $0.\overline{6788}$ (B) $0.\overline{689}$
(C) $0.\overline{6878}$ (D) $0.\overline{687}$

- Q.19** Expressing $0.\overline{23} + 0.\overline{23}$ of a single decimal, we get - [Raj. NTSE Stage-II 2017]

- (A) $0.\overline{465}$ (B) $0.\overline{465}$
(C) $0.\overline{465}$ (D) $0.\overline{4654}$

- Q.20** The value of $\left[\sqrt[3]{\sqrt[6]{a^9}}\right]^4 \left[\sqrt[6]{\sqrt[3]{a^9}}\right]^4$ is :

[NTSE-2012-13]

- (A) a^{16} (B) a^{12} (C) a^8 (D) a^4

- Q.21** If $a = \frac{9}{\sqrt{11}-\sqrt{2}}$; $b = \frac{6}{3\sqrt{3}}$, then the relation between a and b is -

[NTSE-2012-13]

- (A) $a < b$ (B) $a > b$
(C) $a + b < 1$ (D) $a = b$

- Q.22** Value of $\frac{2^{100}}{2}$ is - [NTSE-2013]

- (A) 1 (B) 50^{100} (C) 2^{50} (D) 2^{99}

- Q.23** Which is the greatest number amongst $2^{1/2}$, $3^{1/3}$, $8^{1/8}$ and $9^{1/9}$?

[NTSE Stage-II/2011]

- (A) $9^{1/9}$ (B) $8^{1/8}$ (C) $3^{1/3}$ (D) $2^{1/2}$

- Q.24** If $N = \frac{\sqrt{\sqrt{5}+2} + \sqrt{\sqrt{5}-2}}{\sqrt{\sqrt{5}+2}} - \sqrt{3-2\sqrt{2}}$ then

the value of N is :

[Haryana NTSE Stage-I 2013]

- (A) $2\sqrt{2}-1$ (B) 2
(C) 1 (D) $\sqrt{5}-\sqrt{2}$

- Q.25** If $x^{p^q} = (x^p)^q$, then p =

[Haryana NTSE Stage-I 2013]

- (A) $q^{\frac{1}{q}}$ (B) 1 (C) q^q (D) $q^{\frac{1}{q-1}}$

- Q.26** If in $\sqrt{3} + \sqrt[3]{5}$, $x = \sqrt{3}$ and $y = \sqrt[3]{5}$, then its rationalising factor is -

[M.P. NTSE Stage-I 2013]

- (A) $x + y$
(B) $x - y$
(C) $x^5 + x^4y + x^3y^2 + x^2y^3 + xy^4 + y^5$
(D) $x^5 - x^4y + x^3y^2 - x^2y^3 + xy^4 - y^5$

- Q.27** Of the following four numbers the largest is : [Harayana NTSE Stage-1 2014]

- (A) 3^{210} (B) 7^{140}
(C) $(17)^{105}$ (D) $(31)^{84}$

- Q.41** If the difference of two numbers is 5 and difference of their squares is 300, then sum of the numbers is -

[Rajasthan NTSE Stage-I 2016]

- (A) 1500 (B) 6
(C) 12 (D) 60

- Q.42** The square of an odd integer must be of the form - **[NTSE Stage-II 2013]**

- (A) $6n + 1$
(B) $6n + 3$
(C) $8n + 1$
(D) $4n + 1$ but may not be $8n + 1$

- Q.43** $\sqrt{(a-b)^2} + \sqrt{(b-a)^2}$ is -

[NTSE Stage-2 2013]

- (A) Always zero
(B) Never zero
(C) Positive if and only if $a > b$
(D) Positive only if $a \geq b$

- Q.44** The number of integers n (< 20) for which $n^2 - 3n + 3$ is a perfect square is

[NTSE Stage-II 2014]

- (A) 0 (B) 1 (C) 2 (D) 3

- Q.45** $\sqrt{m^4 n^4} \times \sqrt[6]{m^2 n^2} \times \sqrt[3]{m^2 n^2} = (mn)^k$, then find the value of k -

[Maharashtra NTSE Stage-I 2017]

- (A) 6 (B) 3 (C) 2 (D) 1

- Q.46** If $\frac{3+2\sqrt{3}}{3-\sqrt{3}} = a + \sqrt{3}b$, then the value of $\sqrt{a+b}$ where a and b are rational numbers is -

[Rajasthan NTSE Stage-I 2018]

- (A) 5 (B) 8 (C) 2 (D) 16

- Q.47** Which is unit digit of $6^{18} - 5^{10}$?

[Rajasthan NTSE Stage-I 2018]

- (A) 5 (B) 8 (C) 1 (D) 9

- Q.48** Which of the following is not an irrational number?

[Rajasthan NTSE Stage-I 2019]

- (A) $2 + \sqrt{5}$ (B) $\sqrt{2}$
(C) $\frac{7}{\sqrt{5}}$ (D) $\frac{2\sqrt{11}}{7\sqrt{11}}$

- Q.49** If G.C.D of two number is 8 and their product is 384, then their L.C.M is _____.

[Gujarat NTSE Stage-I 2019]

- (A) 24 (B) 16 (C) 32 (D) 48

- Q.50** $317/3125$ represents

[Gujarat NTSE Stage-I 2019]

- (A) A terminating decimal
(B) A non-recurring decimal
(C) A recurring decimal
(D) An integer

ANSWER KEY

EXERCISE - 1

1. (i) Rational (ii) Rational (iii) Irrational (iv) Rational (v) Rational (vi) Irrational
(vii) Irrational (viii) Irrational (ix) Irrational (x) Irrational
4. 1679 5. 43 6. $\sqrt{12}$, $(108)^{1/4}$ 7. $\sqrt{5}$, $(20)^{1/4}$ 8. No
11. (i) 0.115 (ii) 0.192 (iii) 0.053125 (iv) 0.21375 (v) 0.009375 (vi) 0.00608
12. (i) $\frac{8}{9}$ (ii) $\frac{22}{9}$ (iii) $\frac{8}{33}$ (iv) $\frac{11}{90}$ (v) $\frac{101}{45}$ (vi) $\frac{181}{495}$
13. (i) HCF = 18, LCM = 1584 (ii) HCF = 36, LCM = 11880 (iii) HCF = 128, LCM = 14976
(iv) HCF = 1, LCM = 1260
14. 207 15. 12600 16. 7 m 17. 35 cm 18. 14 19. 9.6 m 20. 8 : 7 : 12 a.m.
21. 10 : 31 hrs 22. (i) 45 (ii) 36 (iii) 15 25. 814 26. 16 times
28. $x = -38$, $y = 13$ 29. (a) 2520 cm 30. (a) 17 l

EXERCISE - 2

Ques.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
Ans.	C	B	C	D	C	B	D	B	A	D	C	C	B	B	A	C	D	A	C	C
Ques.	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40
Ans.	A	C	D	D	D	D	B	B	B	D	D	C	B	D	D	D	A	B	D	A
Ques.	41	42	43	44	45	46	47	48	49	50										
Ans.	A	A	B	B	A	A	C	B	B	A										

EXERCISE - 3

Ques.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
Ans.	A	B	C	A	B	C	B	B	A	C	D	B	D	B	C	D	A	D	B	D
Ques.	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40
Ans.	B	D	C	D	D	D	C	B	C	D	C	D	C	D	C	B	C	B	A	C
Ques.	41	42	43	44	45	46	47	48	49	50										
Ans.	D	C	D	C	B	C	C	D	D	A										

LOGICAL SEQUENCE OF WORDS

In this type of question, some words are given. You have to arrange these words in a meaningful order. The order may be according to age, size and need etc..

Some common sequences have been discussed below :

◆ **TYPE –I : Sequence of occurrence of events or various stages in a process :**

Ex.1 Arrange the following in a meaningful sequence :

1. Leaf 2. Fruit 3. Stem
4. Root 5. Flower

- (A) 4, 3, 1, 5, 2 (B) 2, 3, 4, 1, 5
(C) 4, 3, 1, 2, 5 (D) 5, 1, 4, 3, 2

Sol. (A) Start from **root**, then **stem** will come out and then **leaf**, then **flower** and finally **fruit** will come..

Ex.2 Arrange the following in a logical order :

1. Birth 2. Death
3. Funeral 4. Marriage
5. Education

- (A) 1, 3, 4, 5, 2 (B) 1, 5, 4, 2, 3
(C) 2, 3, 4, 5, 1 (D) 4, 5, 3, 1, 2

Sol. (B) Clearly, the given words when arranged in the order of various events as they occur in a man's life, form the sequence : Birth, Education, Marriage, Death, Funeral.

◆ **TYPE –II : Sequence of objects in a class or group, from part to the whole :**

Ex.3 Arrange the following in a meaningful order, from particular to general:

1. Nation 2. Colony 3. City
4. District 5. State

- (A) 2, 3, 4, 5, 1 (B) 3, 1, 2, 5, 4
(C) 3, 1, 4, 2, 5 (D) 3, 1, 4, 5, 2

Sol. (A) Clearly, a **Colony** is a part of a **City**, which in turn is a part of **District** which is part of a **State** which lies within a **Country**.

Ex.4 Arrange the following in a logical order :

1. Shoulder 2. Wrist 3. Elbow
4. Palm 5. Finger

- (A) 2, 4, 5, 3, 1 (B) 3, 1, 4, 2, 5
(C) 3, 4, 5, 2, 1 (D) 5, 4, 2, 3, 1

Sol. (D) Clearly, we are given the names of parts of a hand, which may be arranged

(i) from top to bottom, i.e., **Shoulder, Elbow, Wrist, Palm, Finger**, which is 1, 3, 2, 4, 5; or

(ii) from bottom to top, i.e., **Finger, Palm, Wrist, Elbow, Shoulder**, which is 5, 4, 2, 3, 1.

Out of these, the sequence 5, 4, 2, 3, 1 is given in the alternatives provided.

◆ **TYPE –III : Sequence of increasing /decreasing size, value, intensity etc. :**

Ex.5 Arrange the following in a logical sequence from small to big :

1. Elephant 2. Cat 3. Mosquito
4. Tiger 5. Whale

- (A) 2, 3, 1, 4, 5 (B) 3, 2, 4, 1, 5
(C) 5, 1, 4, 2, 3 (D) 5, 4, 1, 2, 3

Sol. (C) Clearly, The **Whale** is the biggest then smaller than whale is **Elephant** then **Tiger** then **Cat** and then the smallest is **Mosquito**.

Ex.6 Arrange the following in a logical order :

1. Gold 2. Iron 3. Sand
4. Platinum 5. Diamond

- (A) 2, 4, 3, 5, 1 (B) 3, 2, 1, 4, 5
(C) 4, 5, 1, 3, 2 (D) 5, 4, 3, 2, 1

Sol. (B) Clearly, the given names when arranged in order of increasing values, i.e., from cheapest to the most expensive, form the sequence: **Sand, Iron, Gold, Platinum, Diamond**.

Thus, the correct answer is 3, 2, 1, 5, 4.

EXERCISE

Directions : (Q.1 to Q.30) : In each of the following questions, arrange the given words in a meaningful sequence and then choose the most appropriate sequence from amongst the alternatives provided below each question :

- Q.1** 1. Skull 2. Shoulder
3. Neck 4. Face
5. Legs
(A) 1, 2, 3, 4, 5 (B) 1, 4, 3, 2, 5
(C) 1, 3, 4, 2, 5 (D) 1, 4, 2, 3, 5

- Q.2** 1. Sun 2. Planet
3. Galaxy 4. Stars
5. Moon
(A) 1, 3, 2, 5, 4 (B) 1, 3, 2, 4, 5
(C) 3, 1, 2, 5, 4 (D) 3, 4, 1, 2, 5

- Q.3** 1. Teacher 2. College
3. Guide 4. Study
5. Exam
(A) 2, 3, 1, 4, 5 (B) 2, 4, 1, 3, 5
(C) 4, 1, 2, 3, 5 (D) 2, 1, 3, 4, 5

- Q.4** 1. Animals 2. Cow
3. Vertebrates 4. Mammals
(A) 4, 1, 2, 3 (B) 1, 4, 2, 3
(C) 1, 3, 4, 2 (D) 4, 1, 2, 3

- Q.5** 1. Flat 2. Home
3. Building 4. Society
5. Place
(A) 5, 4, 3, 1, 2 (B) 5, 4, 3, 2, 1
(C) 4, 3, 1, 2, 5 (D) 4, 5, 3, 1, 2

- Q.6** 1. Sentence 2. Chapter
3. Letter 4. Book
5. Word 6. Paragraph
(A) 4, 2, 1, 6, 5, 3 (B) 4, 2, 6, 1, 5, 3
(C) 4, 6, 1, 2, 3, 5 (D) 4, 6, 2, 5, 1, 3

- Q.7** 1. Cut 2. Put on
3. Mark 4. Measure
5. Tailor
(A) 1, 3, 2, 4, 5 (B) 2, 4, 3, 1, 5
(C) 3, 1, 5, 4, 2 (D) 4, 3, 1, 5, 2

- Q.8** 1. Country 2. Furniture
3. Forest 4. Wood
5. Trees
(A) 1, 3, 5, 4, 2 (B) 1, 4, 3, 2, 5
(C) 2, 4, 3, 1, 5 (D) 5, 2, 3, 1, 4

- Q.9** 1. Education 2. Job
3. Selection 4. Salary
5. Interview
(A) 1, 3, 5, 2, 4 (B) 1, 5, 3, 2, 4
(C) 1, 2, 3, 5, 4 (D) 1, 5, 2, 4, 3

- Q.10** 1. Child 2. Play 3. Game 4. Chess
5. Happy
(A) 1, 3, 2, 4, 5 (B) 1, 2, 3, 4, 5
(C) 1, 3, 4, 2, 5 (D) 5, 1, 2, 3, 4

- Q.11** 1. Index 2. Chapter
3. Book 4. Introduction
5. Questions
(A) 3, 4, 1, 2, 5 (B) 3, 1, 4, 2, 5
(C) 3, 1, 2, 5, 4 (D) 4, 3, 1, 2, 5

- Q.12** 1. Grain 2. Plant
3. Sandwich 4. Bread
5. Dough
(A) 1, 2, 5, 4, 3 (B) 2, 1, 4, 5, 3
(C) 2, 1, 5, 4, 3 (D) 2, 1, 4, 5, 3

- Q.13** 1. Andhra Pradesh 2. Universe
3. Tirupati 4. World
5. India
(A) 1, 5, 3, 2, 4 (B) 2, 1, 3, 5, 4
(C) 3, 1, 5, 4, 2 (D) 5, 4, 2, 1, 3

- Q.14** 1. Atomic Age 2. Metallic Age
3. Stone Age 4. Alloy Age
(A) 1, 3, 4, 2 (B) 2, 3, 1, 4
(C) 3, 2, 4, 1 (D) 4, 3, 2, 1

- Q.15** 1. Office 2. Save
3. Work 4. Earn
5. Spend
(A) 1, 3, 4, 2, 5 (B) 1, 4, 3, 5, 2
(C) 1, 3, 4, 5, 2 (D) 3, 1, 4, 5, 2

- Q.16** 1. Planet 2. Continents
3. Countries 4. Cities
5. Land
(A) 1,2,5,3,4 (B) 1,5,2,3,4
(C) 1,2,3,5,4 (D) 1,3,4,2,5
- Q.17** 1. Post-box 2. Letter
3. Envelope 4. Delivery
5. Clearance
(A) 2, 3, 1, 4, 5 (B) 3, 2, 1, 4, 5
(C) 3, 2, 1, 5, 4 (D) 3, 2, 4, 5, 1
- Q.18** 1. Foetus 2. Child
3. Baby 4. Adult
5. Youth
(A) 1, 2, 4, 3, 5 (B) 1, 3, 2, 5, 4
(C) 2, 3, 5, 4, 1 (D) 5, 4, 2, 3, 1
- Q.19** 1. Poverty 2. Population
3. Death 4. Unemployment
5. Disease
(A) 1,2,3,4,5 (B) 2,4,1,5,3
(C) 3,4,2,5,1 (D) 2,3,4,5,1
- Q.20** 1. Heel 2. Shoulder
3. Skull 4. Neck
5. Knee 6. Chest
7. Thigh 8. Stomach
9. Face 10. Hand
(A) 2, 4, 7, 10, 1, 5, 8, 9, 6, 3
(B) 3, 4, 7, 9, 2, 5, 8, 10, 6, 1
(C) 4, 7, 10, 1, 9, 6, 3, 2, 5, 8
(D) 3, 9, 4, 2, 10, 6, 8, 7, 5, 1
- Q.21** 1. Rain 2. Monsoon
3. Rescue 4. Flood
5. Shelter 6. Relief
(A) 1, 2, 3, 4, 5, 6 (B) 1, 2, 4, 5, 3, 6
(C) 2, 1, 4, 3, 5, 6 (D) 4, 1, 2, 3, 5, 6
- Q.22** 1. Farm 2. Flowers
3. Trees 4. Fruits
5. Juice
(A) 1,2,3,4,5 (B) 3,1,4,2,5
(C) 1,3,2,4,5 (D) 3,1,2,4,5

- Q.23** 1. Ceiling 2. Room
3. Floor 4. Walls
5. Foundation
(A) 5, 4, 1, 3, 2 (B) 5, 4, 3, 1, 2
(C) 4, 5, 3, 1, 2 (D) 4, 5, 1, 2, 3
- Q.24** 1. Puberty 2. Adulthood
3. Childhood 4. Infancy
5. Senescence 6. Adolescence
(A) 2, 4, 6, 3, 1, 5 (B) 4, 3, 1, 6, 2, 5
(C) 4, 3, 6, 2, 1, 5 (D) 5, 6, 2, 3, 4, 1
- Q.25** 1. Cutting 2. Dish
3. Vegetable 4. Market
5. Cooking
(A) 1, 2, 4, 5, 3 (B) 3, 2, 5, 1, 4
(C) 4, 3, 1, 5, 2 (D) 5, 3, 2, 1, 4
- Q.26** 1. Never 2. Sometimes
3. Generally 4. Seldom
5. Always
(A) 5, 2, 1, 3, 4 (B) 5, 2, 4, 3, 1
(C) 5, 3, 2, 1, 4 (D) 5, 3, 2, 4, 1
- Q.27** 1. Table 2. Tree
3. Wood 4. Seed
5. Plant
(A) 1, 2, 3, 4, 5 (B) 1, 3, 2, 4, 5
(C) 4, 5, 2, 3, 1 (D) 4, 5, 3, 2, 1
- Q.28** 1. Milky way 2. Sun
3. Moon 4. Earth
5. Stars
(A) 1, 4, 3, 2, 5 (B) 2, 3, 4, 5, 1
(C) 3, 4, 2, 5, 1 (D) 4, 3, 2, 5, 1
- Q.29** 1. Sea 2. Rivulet
3. Ocean 4. River
5. Glacier
(A) 5, 2, 1, 3, 4 (B) 5, 2, 4, 1, 3
(C) 5, 4, 2, 3, 1 (D) 5, 4, 3, 2, 1
- Q.30** 1. Plane 2. Take off
3. Land 4. Airport
5. Runway
(A) 4, 1, 5, 3, 2 (B) 4, 5, 3, 1, 2
(C) 4, 5, 2, 1, 3 (D) 4, 5, 1, 2, 3

Q.31 Arrange the following in a meaningful sequence:
A = Birth, B = Death, C = Funeral, D = Marriage, E = Education.

[Rajasthan NTSE Stage-I 2019]

- (A) AEDBC (B) ADECB
(C) AEBDC (D) ADEBC

Q.32 Arrange the following in a meaningful sequence:
A = Study, B = Service, C = Examination, D = Earning, E = Result.

[Rajasthan NTSE Stage-I 2019]

- (A) EACDB (B) ABECD
(C) ACEBD (D) AECBD

Q.33 Arrange the following words in the sequence in which they occur in the dictionary, then choose the correct option

[Delhi NTSE Stage-I 2019]

- (i) BHAGWAN (ii) BHAGWAT
(iii) BHAGIRATH (iv) BHAGAT
(A) iv, i, iii, ii (B) iv, ii, i, iii
(C) iv, iii, ii, i (D) iv, iii, i, ii

Instruction: According to the question, Choose the correct option logically in question no. 34 to 37.

[Gujarat NTSE Stage-I 2019]

- Q.34** (1) Embryo (2) Child
(3) Baby (4) Middle Aged
(5) Young
(A) 1,3,4,5,2 (B) 1,3,5,2,4
(C) 1,3,2,5,4 (D) 1,3,4,2,5

- Q.35** (1) Poverty
(2) Population
(3) death
(4) Unemployment
(5) Disease
(A) 2,4,1,5,3 (B) 1,2,3,4,5
(C) 2,3,4,5,1 (D) 2,4,5,1,3

- Q.36** (1) Accident (2) Judge
(3) Doctor (4) Lawyer
(5) Police
(A) 1,3,4,2,5 (B) 1,3,5,4,2
(C) 1,2,3,4,5 (D) 1,2,5,4,3

- Q.37** (1) Golden Jubilee
(2) Silver Jubilee
(3) Anniversary
(4) Diamond Jubilee
(5) Centenary Celebrations
(A) 2,1,3,4,5 (B) 2,3,4,5,1
(C) 3,1,2,4,5 (D) 3,2,1,4,5

THE RISE OF NATIONALISM IN EUROPE

Chapter Outline

- ✧ The Theme of Frederic Sorrieu's Painting
- ✧ French Revolution and the idea of the nation
- ✧ The Making of Nationalism in Europe
- ✧ The age of Revolutions 1830-1848
- ✧ The Making of Germany and Italy
- ✧ Visualising the Nation
- ✧ Nationalism and Imperialism

The Theme of Frederic Sorrieu's Painting



The Dream of Worldwide Democratic and Social Republics - The Pact Between Nations, a print prepared by Frederic Sorrieu, 1848.

- (i) In 1848, Frederic Sorrieu, a French artist, prepared a series of four prints visualizing his dream of a world made up of 'democratic and social Republics', as he called them.
- (ii) The first print of the series, shows the people of Europe and America – men and women of all ages and social classes – marching in a long train and offering homage to the statue of Liberty as they pass by it.

- (ii) A female figure was shown with the torch of Enlightenment in one hand and the Charter of the Rights of Man in the other.
- (iii) On the earth in the foreground of the image lie the shattered remains of the symbols of absolutist institutions.
- (iv) In Sorrieu's utopian vision, the peoples of the world are grouped as distinct nations, identified through their flags and national costume.

The French Revolution and The Idea of the Nation

France was a full fledged territorial state in 1789 under the rule of an absolute monarch. The political and constitutional changes that came in the wake of French Revolution led to the transfer of sovereignty from the monarchy to a body of French citizens. The **revolution proclaimed** that it was the people who would hence forth constitute the nation and shape its destiny.

◆ Steps Taken by French Revolutionaries :

- (i) The ideas of **La Patrie (the fatherland)** and **Le Citoyen (the citizen)** emphasized the notion of a united community enjoying equal rights under a constitution.
- (ii) A new French **flag, the tricolour**, was chosen to replace the former royal standard. New **hymns** were composed, **oaths** taken and martyrs commemorated, all in the name of a **nation**.
- (iii) The Estates General was elected by the body of active citizens and renamed the **National Assembly**.
- (iv) A **centralised administrative system** was put in place and it formulated **uniform laws** for all citizens within its territory.
- (v) Internal customs duties and dues were abolished and a **uniform system of weights and measures** was adopted.
- (vi) Regional dialects were discouraged and French, as it was spoken and written in Paris, became the common language of the nation.
- (vii) The revolutionaries further declared that it is was the mission and the destiny of the French nation to liberate the peoples of Europe from despotism.

◆ Civil Code of 1804 :

- (i) Though a return to monarchy Napoleon had, no doubt, destroyed democracy in France, but in the administrative field he had incorporated revolutionary principles in order to make the whole system more rational and efficient.
- (ii) The civil code of 1804-usually known as the **Napoleonic code** did away with all privileges based on birth, established equality before the law and secured the right to property.
- (iii) This code was exported to the regions under French Control.
- (iv) In the Dutch Republic in Switzerland, in Italy and Germany. Napoleon abolished the feudal system and freed peasants from serfdom and manorial dues.

- (v) In the towns too guild restrictions were removed.
- (vi) Transport and communication systems were improved
- (vii) Peasants, artisans, workers and new businessmen enjoyed a new-found freedom.



The Courier of Rhineland loses all that he has on his way home from, Leipzig. Napoleon here is represented as a postman on his way back to France after he lost the battle of Leipzig in 1813. Each letter dropping out of his bag bears the names of the territories he lost.

The Making of Nationalism in Europe

In the mid-eighteenth century, Germany, Italy and Switzerland were divided into Kingdoms, duchies and cantons whose rulers had their autonomous territories. Eastern and Central Europe were under autocratic monarchies within the territories of which lived diverse people. They did not see themselves as Sharing a collective identity or a common culture. Such differences did not easily promote a sense of political unity. The only tie binding these diverse groups together was common allegiance to the emperor.

COMPETITIVE LEVEL

- 1797 Napoleon invades Italy; Napoleonic wars begin.
- 1814-1815 Fall of Napoleon; the Vienna Peace Settlement.
- 1821 Greek struggle for independence begins.
- 1848 Revolutions in Europe; artisans, industrial workers and peasants revolt against economic hardships ; middle classes demand constitutions and representative governments; Italians, Germans, Magyars, Poles, Czechs, etc. demand nation-states.
- 1859-1870 Unification of Italy.
- 1866-1871 Unification of Germany. 1905 Slav nationalism gathers force in the Habsburg and Ottoman Empires.



(i) The Aristocracy and the New Middle Class :

- (1) The members of this class were united by a common way of life that cut across regional diversions. They owned castles in the countryside and also town houses. This powerful autocracy was however, numerically a small group. The majority of the population was made up of the peasantry.

GLOSSARY

1. **Nationalism** : A sense of belonging to one nation. Feeling of pride and patriotism towards the country one belongs.
2. **Absolutist** : Unrestricted, despotic and authoritarian often refers to a centralised repressive monarchical government.
3. **Utopian** : An ideal situation a vision too good to be realised in practice.
4. **Nation state** : A state having a common and contiguous boundary with inhabitants, people sharing common language, race and religion. Majority of its citizen develop a sense of common identity and share a common history.
5. **Plebiscite** : A direct vote by which the people of a region, themselves decide to accept or reject a proposal.
6. **Sovereignty** : Supreme Power.
7. **Monarchy** : Form a government headed by a monarch or a hereditary or dynastic ruler.
8. **The Estates General** : Referred to the French Parliament an elected body which was renamed as the National Assembly after the revolution of 1789.
9. **Civil Code** : A systematic set of laws for the citizen.
10. **Habsburg Empire** : The empire that ruled Austria, Hungary including the Alpine region of Tyrol, Austria, Sudetenland, Bohemia. It also included Italian provinces of Lombardy and Venetia.
11. **Liberalism** : Derived from the word 'liber' meaning free. The idea of liberalism stands for freedom of individual and equality of all before law. Politically it refers to representative government.
12. **Suffrage** : The right to vote.
13. **Elle** : Elle was used to measure cloth, prevalent in German states.
14. **Zollverein** : A custom union, formed in 1834 in Prussia to remove barriers of trade.
15. **Conservatism** : A spirit or philosophy which believes in maintaining and preserving traditional values and institutions. It prefers gradual change to quick and drastic change.
16. **Carbonari** : A secret society of Italy consisting of young revolutionaries.
17. **Young Italy** : A secret society founded by Mazzini at Masses for organising revolutionary activities.
18. **Ottoman Empire** : Turkish empire ruled by the caliph. The spiritual and temporal head of the muslim.
19. **Romanticism** : A cultural movement which aimed at developing a particular form of national sentiment and promote a feeling of collective heritage as the basis of motion.
20. **Das volk** : A German word meaning common people.
21. **Republic** : A state where the head of the state is elected and does not hold a hereditary position.
22. **Feminist** : People who advocate women's right on the basis of equality of sexes.
23. **Ideology** : System of ideas reflecting a particular social and political vision.
24. **Allegory** : Symbol representing an abstract idea; an idea identified through a person or a thing.
25. **Balkan region** : A region in Europe with geographical and ethnic variation. The region covers the states of modern day Romania, Bulgaria, Albania, Greece, Macedonia, Croatia, Bosnia, Herzegovina, Slovenia, Serbia, Montenegro. The inhabitants of the region are known as the Slavs.

EXERCISE-1

➤ Very Short Answer Type Questions

- Q.1** What is “Statue of Liberty”? What does the statue signify ?
- Q.2** Name four European powers who collectively defeated Napoleon.
- Q.3** With what aim was the Treaty of Vienna (1815) signed?
- Q.4** When and under whom was the constitutional monarchy setup in France?
- Q.5** Name the treaty which recognised Greece as an independent nation.

➤ Short Answer Type Questions

- Q.6** What are the main symbols of an independent nation ?
- Q.7** Examine any four reasons for the nationalistic upsurge in 19th century Europe ?
- Q.8** What are the most important achievements of the French Revolution of 1789 ?
- Q.9** What was the significance of the Napoleonic code ?
- Q.10** Examine the reasons for the political disunity in Habsburg empire of Austria and Hungary.
- Q.11** How did the growth of industrialisation change the social and political equation of Europe ?
- Q.12** Examine the liberal ideology imbibed by the revolutionary leader after French revolutions.
- Q.13** How did liberalism give rise to economic nationalism?

Q.14 What was the impact of Treaty of Vienna (1815) on European people?

Q.15 Describe the contribution of Mazzini in the unification of Italy?

➤ Long Answer Type Questions

Q.16 Mention the main factors responsible for the rise of nation state.

Q.17 What were the major proposals of the Vienna Congress.

Q.18 Briefly trace the process of the unification of Germany and that of Britain.

Q.19 Describe any features of the Vienna Treaty of 1815 in brief.

Q.20 Who was Count Camillo de Cavour? Mention his contributions.

EXERCISE-2

- Q.1** A nation state, according to the French philosopher Ernst Renan, was formed by
(A) A common language, race, religion or territory
(B) A nation with a shared history or descent, of endeavors, sacrifice and devotion, wish to perform still more great deeds together
(C) A large scale solidarity, its existence a daily plebiscite
(D) Both (B) and (C)
- Q.2** Federic Serrieu, a French artist, in his series of four prints (1848) visualised his dream of a world as :
(A) A world made up of 'democratic and social republics'
(B) A world made up of one nation, one world
(C) A world with one absolute ruler
(D) A world following one religion, one language
- Q.3** The Civil Code of 1804, also known as the Napoleonic Code, established :
(A) Equality before the law
(B) Secured the right to property
(C) Did away with all the privileges based on birth
(D) All the above
- Q.4** Which new spirit guided European nations after Napoleon's defeat?
(A) Fascism (B) Conservatism
(C) Nazism (D) Communism
- Q.5** All the new regimes, set up in 1815, were autocratic because :
(A) They did not tolerate criticism and dissent
(B) They imposed censorship laws to control what was said in newspapers, plays, songs etc.
(C) They curbed activities which questioned their legitimacy
(D) All the above
- Q.6** The purpose behind the painting "The Massacre at Chios" by Eugene Delacroix, 1824, was:
(A) To appeal to the emotions of the spectators and create sympathy for the Greeks
(B) To dramatise the incident in which 20,000 Greeks were killed
(C) To focus on the suffering of women and children
(D) All the above
- Q.7** German philosopher Johann Gottfried Herder claimed that true German culture was discovered through
(A) Folk songs, folk poetry, folk dances
(B) Common people - das volk
(C) Vernacular language
(D) All the above
- Q.8** The weavers of Silesia revolted in 1845 against contractors because :
(A) The contractors who gave them orders drastically reduced their payments
(B) The contractors took advantage of their misery and desperate need for jobs
(C) Both (A) and (B)
(D) The contractors had killed eleven weavers
- Q.9** The artists of the 18th and 19th centuries personified a nation as :
(A) A particular woman to represent the nation
(B) A female figure
(C) A female figure as an allegory, to represent an abstract idea of a nation in concrete form
(D) All the above

- Q.10** Phillip Veit used the following in his painting as symbols
 (i) Broken chains,
 (ii) Sword
 (iii) Olive branch round the sword and
 (iv) Rays of the rising Sun.
 They symbolised :
 (A) (i) Heroism,
 (ii) readiness to fight
 (iii) strength
 (iv) hope
 (B) (i) Freedom
 (ii) strength
 (iii) readiness to fight
 (iv) Beginning of a new era
 (C) (i) Freedom
 (ii) readiness to fight
 (iii) willingness to make peace
 (iv) beginning of a new era
 (D) (i) Heroism
 (ii) courage
 (iii) readiness to fight
 (iv) hope
- Q.11** The two reasons which made the Balkans an explosive region after 1871 were :
 (A) The spread of ideas of romantic nationalism and the disintegration of the Ottoman Empire under whose control they were
 (B) The declaration of independence by European subject nationalities from the Ottoman Empire.
 (C) The support of history and nationalism to be independent
 (D) All the above
- Q.12** Returning from exile, Giuseppe Mazzini formed a new organisation called
 (A) National Italy (B) Young Italy
 (C) United Italy (D) Organised Italy
- Q.13** System of ideas reflecting a particular social and political vision is known as
 (A) Ideology (B) Pedagogy
 (C) Philology (D) Genealogy
- Q.14** Guiseppe Garibaldi led an army of volunteers to Rome to fight the last obstacle in the unification of Italy in the year
 (A) 1857 (B) 1867
 (C) 1877 (D) 1887
- Q.15** Napoleon invaded Italy in the year
 (A) 1781 (B) 1782
 (C) 1787 (D) 1789
- Q.16** An abstract idea or emotion which when used as a symbol to portray a theme with respect to a nation is called
 (A) An ideology (B) symbol
 (C) an allegory (D) a painted theme
- Q.17** The word das volk refers to
 (A) common people of France
 (B) common people of Italy
 (C) common people of Germany
 (D) common people of Russia
- Q.18** The theory that tries to make awareness of women's rights and interests based on the belief of the social, economic and political equality of the genders is known as
 (A) Humanism (B) Feminism
 (C) Post modernism (D) Culturalism
- Q.19** la patrie, one of the ideas used during the French Revolution to emphasize the notion of a united community, means
 (A) Holy land (B) Fatherland
 (C) Motherland (D) United land
- Q.20** The term 'liberalism' is derived from
 (A) Latin language
 (B) Russian language
 (C) French language
 (D) Greek language
- Q.21** The Architect of German unification process was
 (A) Otto von Bismarck
 (B) Hitler
 (C) Giuseppe Mazzini
 (D) John Gottfried Herder
- Q.22** In Ireland a revolt by Catholic Irishmen in the year 1798 was led by
 (A) Milton Booth (B) Wolfe Tone
 (C) McGregor (D) Potemkin
- Q.23** The Grimms brothers were
 (A) British nationals
 (B) French nationals
 (C) German nationals
 (D) Italian nationals

EXERCISE-3

- Q.1** What is the meaning of this French word "Le Citoyen" ? [Haryana-NTSE Stage-1/2013]
 (A) The people (B) The Citizen
 (C) Resident (D) All above
- Q.2** Chief Minister Cavour who led the movement of unification of Italy was a [Haryana-NTSE Stage-1/2013]
 (A) Freedom Fighter
 (B) A revolutionary
 (C) A democrat
 (D) Neither a Revolutionary nor a Democrat
- Q.3** Who hosted Congress of Vienna ? [Punjab-NTSE Stage-1/2013]
 (A) Bismark
 (B) Napoleon Bonaparte
 (C) Voltaire
 (D) Malternich
- Q.4** Who is regarded as father of Italian unification ? [M.P.-NTSE Stage-1/2013]
 (A) Mazini (B) Cavour
 (C) Garibaldi (D) None of these
- Q.5** Who amongst the following ruled over Sardinia-pied mont during the middle of the nineteenth century. [Haryana-NTSE Stage-1/2013-14]
 (A) Italian Princely house
 (B) Austrian Habsburg
 (C) Pope
 (D) Bourbon king of Spain
- Q.6** Who was Paul Bernard? [Haryana-NTSE Stage-1/2013-14]
 (A) Capitalist (B) Social worker
 (C) Social reformer (D) Economist
- Q.7** Who said "When France sneezes the rest of the Europe catches cold". [Haryana-NTSE Stage-1/2013-14]
 (A) Garibaldi (B) Mazzini
 (C) Bismarck (D) Matternich
- Q.8** Find out the statement which does not cause to Imperialism : [Maharashtra-NTSE Stage-1/2014]
 (A) Prosperity of Asia and Africa
 (B) Weakness of Asian and African Nations
 (C) Need of raw material from Asia and Africa
 (D) Growing agitation in Nationalistic movement in African and Asian continent
- Q.9** Germany was unified in - [Haryana-NTSE Stage-1/2013-14]
 (A) 1870 (B) 1871
 (C) 1872 (D) 1873
- Q.10** The tactful diplomatic alliance between Sardinia-Piedmont and France was engineered by [Rajasthan-NTSE Stage-1/2015]
 (A) Mazzini
 (B) Cavour
 (C) Garibaldi
 (D) Victor Emmanuel
- Q.11** Which of following state was ruled by as Italian Princely House ? [Haryana-NTSE Stage-1/2015]
 (A) Sardia Piedmont
 (B) Papal States
 (C) Venetia
 (D) Tuscany
- Q.12** How many years did it take to form Republic of Vietnam ? [Punjab-NTSE-Stage-1-2015]
 (A) 30 (B) 29
 (C) 19 (D) 25
- Q.13** Which French artist prepared a series of four prints visulizing his dream of a world made up of 'Democratic and Social Republics'? [Punjab-NTSE-Stage-1-2015]
 (A) Frederic Sorrieu
 (B) Andreas Rebmann
 (C) Karl Kaspar Fritz
 (D) Giuseppe Mazzini

ANSWER KEY

EXERCISE - 2

Ques.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Ans.	D	A	D	B	D	D	D	C	C	D	A	B	A	B	C
Ques.	16	17	18	19	20	21	22	23							
Ans.	C	C	B	B	A	A	B	B							

EXERCISE - 3

Ques.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Ans.	B	D	D	B	A	D	D	D	B	B	A	A	A	A	A
Ques.	16														
Ans.	B														

LETTER WRITING

Formal Letter

◆ LETTER TO EDITOR

Purpose of the Letter

- To express one's opinion on a wide range of social issues.
- To express and share one's observation and views in a public forum, namely a magazine or a newspaper.

Body of the letter

- The presentation should be concise.
- It should clearly establish the cause-effect relationship.
- Efforts taken to rectify the cause of the issue must be stated.

Language of the Letter

- Formal language must be used.
- Repetition should be avoided.
- Sentences should not be very long.
- The letter should be divided into two or three paragraphs.

Important Points

- The letter must be written in the appropriate format.
- The word limit of 150 words must be kept in mind.

◆ Format

Sender's Address

Date

Address of addressee

Dear Sir/Madam

Subject : _____

Body of the letter

1. Statement of problem

- (Through the columns of your newspaper)
- (I was to read)
- (With reference to the news report dated I was to read)

2. Causes of problem

- It is alleged
- is due largely to

3. Results of problem

- has led to
- As a result

4. Suggestions

- I feel
- In my opinion
- I suggest

Yours faithfully / Yours sincerely
name of sender

◆ Sample Formal Letter

You are Sunil Shetty of 7 Club Road, Hyderabad-13. Write a letter to the Editor of The Hyderabad Times. P.O. Box 350 Hyderabad-1, about the bad quality and inadequate supply of tap water in your locality.

Sender's address (only address—not name)	7 Club Raod Hyderabad	April 12, 2011	Date
Designation of receiver	The Editor		
Receiver's address	The Hyderabad Times P.O. Box 350 Hyderabad-1		
Salutation	Sir		
Subject statement	Subject : Inadequate Supply of Water		
Problem	Through the columns of your newspaper, I would like to draw the attention of the civic authorities of the bad quality and inadequate supply of tap water in our locality.		
Effect of problem	It is unfortunate that the civic authorities do not take care to provide the citizens with even the basic amenities such as proper and adequate water supply. In our locality, water is supplied only for an hour in the morning. For the rest of the day, the taps remains dry. Also, the quality of water supplied is unhygienic. there have been some cases of diarrhea and dysentery. It is feared that these diseased may break out in epidemic form if the quality of water is not improved.		
Desired future course of action	It is hoped that instead of ignoring the matter, the authorities concerned will take immediate steps to ensure adequate supply of good quality tap water in our locality.		
Subscription and signature	Yours faithfully Sunil Shetty		

1. Write a letter in 100-120 words to the editor of The Daily Express, expressing your views on the topic, Nukkad Natak. You feel that street play is an effective medium to educate the society about vices which are cancerous. You are Amrit/Amrita, B-94, Malviya Nagar, Kanpur.

Ans. B-94, Malviya Nagar
Kanpur (U.P.)
16th September, 20.....
The Editor
The Daily Express
Lucknow (U.P.)
Sub. : **Importance of nukkad natak**
Sir

Yesterday I watched a nukkad natak in our street and realized that they can play a significant role to educate people about social and political problems. The idea behind the nukkad natak is to educate people through entertainment. The natak I watched was about Swachh Bharat. It highlighted how we can contribute to keeping our neighbourhood clean to remain healthy. It was based on the story of two neighbours. While the behaviour of the two neighbours evoked laughter, the theme was not lost on the audience.

I think nukkad natak can do a lot about educating people about economical use of water, importance of physical exercise, cheap but nutritious diet, planting and looking after trees, kindness to animals, saving money, personal hygiene and so many other things.

The great advantage of the nukkad natak is that it is inexpensive as it is performed in public places such as streets where everyone can watch it without paying admission.

Yours sincerely
Amrita

2. **Children these days are addicted to fast food from the microwave like noodles, pizzas or readymade chips and cookies, instead of the balanced meals they need to take. As Shruti/Sahil, write a letter to the Editor of “The Hindustan Times” in about 100-120 words, expressing your concern over the situation and suggesting measures to combat the problem.**

Ans. 71, Surya Vihar
Ghaziabad (U.P.)
8 June, 20.....
The Editor
Hindustan Times
New Delhi

Sub : Addiction to fast food

Sir

Children these days are getting more and more addicted to fast food. In fact the television has played a great role in introducing fast foods and promoting it in our homes. Big companies pay hefty fees to popular who are the icons of the youth to promote their products. The companies make tall but unverified claims. They promise that their product will make children strong and help them grow faster and make them champions. They also use ambiguous language to misguide people and innocent children who are their easy targets. Mothers too are influenced by audio-video clips and serve such foods to their children. They seldom realize that the people who promote those products seldom use them.

The housewives these days have little time to cook. They are happy to serve instant food without realizing that the food does not only lack nutrients but is also injurious to health. They contain preservatives and other chemicals to make food tastier. Potato chips, noodles, fruit drinks are not what they claim to be. We must revert to our home made food which is hygienic, nutritious, fresh and free from harmful additives.

Yours sincerely

Shruti

3. **Write a letter in 100-120 words, to the editor of The Daily Express expressing your views on the topic “It’s the individuals outlook and the self driven motive that will bring about the Swachh Bharat initiative true and not any forced government campaign.” You are Amrit/ Amrita, B-94, Malviya Nagar Kanpur.**

Ans. B-94, Malviya Nagar
Kanpur (U.P.)
17 June, 20.....
The Editor
The Daily Express
Kanpur (U.P.)

Sub : Swachh Bharat Campaign

Sir

Our honourable Prime Minister wielded a broom himself to give a sweeping start to the Swachh Bharat Campaign. It is a noble idea because cleanliness is next to godliness Gandhiji always laid stress on cleanliness and in the Sabarmati Ashram everybody was required to clean the toilets.

No doubt the government has an important role to play to make the campaign a success. They have to provide bins where people can deposit their garbage. Further, the garbage needs to be collected and disposed off in a proper manner. But it is the people and people alone, who can do a lot. First of all they must see that their cleanliness does not end at their doorstep. They must keep their garbage at proper places. They should not push it on to the street or towards the neighbour's door.

People eat fruit or some other eatables. They litter empty boxes, bottles, polythene bags, fruit peels, etc. everywhere – in the parks, in the buses, trains, streets and elsewhere. Some people spit betel-leaf juice on walls and nooks of public places. These are unhealthy habits and must be curbed. Children can be taught cleanliness both at home and school. People can be educated through television, Police should warn people against spreading litter. Anybody who makes public places dirty, must be made to clean it.

Yours sincerely

Amrita

4. **Science and technology has revolutionised life. It has developed and improved the quality of life in various spheres such as health, environment, agriculture, etc. Using your own ideas and ideas from the unit, 'Science' of MCB, write a letter to the editor of a national daily in about 100-120 words about the improvements due to science and technology.**

Ans. ABC Building

Delhi

October 14, 20.....

The Editor

The Times of India

Delhi

Sub. : **Improvements brought by science and technology**

Sir

Through the columns of your esteemed newspaper, I would like to highlight the improvements and developments in our lives due to science and technology. Science has revolutionised our lives. It has developed and improved the quality of life in various spheres such as health, environment, agriculture, etc. Science has made our lives easy. There are mobile phones, motor cars, aeroplanes, tractors, well equipped medical machines, and tools. All these developments in the field of science have served to save our time, energy and efforts. Further medical research to find a cure for AIDS, modern health care and computers have improved the quality of life.


Finally, I would like to conclude by saying that science and technology do not make undesirable changes to lifestyles, rather they upgrade it.

Thanking you.

Yours sincerely

Abhishek

5. Himanshu Jain is an active social worker. One day he came across a sight which is shown alongside. It touched him deeply. Imagining yourself to be Himanshu write a letter to the editor of a national daily voicing your protest against child labour in about 100-120 words.



Ans. 123, Vasant Building

Mumbai

August 21, 20.....

The Editor

The Tribune

Mumbai

Sub. : Protest against child labour

Sir

Through the columns of your esteemed newspaper, I would like to highlight the problem of child labour in our society. It is really a shame for our society that children who are meant to be the future of our nation, should suffer such hardships. Poor children employed in the industries suffer various abuses. People are picking them for household chores like cleaning and sweeping. They are also engaged in many hazardous jobs. They are deprived of the innocent and simple pleasures of childhood.

It is the responsibility of the government, society and parents to come forward to solve this problem. Children should be provided with free and compulsory elementary education. Child labour has been banned in India. The provisions made by the Constitution in this regard should be strictly implemented. Top priority should be given to projects related to child welfare issues. With these provisions, their problems can be minimised.

Thanking you.

Yours sincerely

Himanshu Jain

EXERCISE

1. You are Suman Sinha of 13, Dilshad Garden, Delhi. You have observed that the subways in Delhi are seldom used by pedestrians. You have decided to write a letter to the Editor of a national daily, highlighting the dire need of creating awareness about it. Based on the points given and ideas from the unit 'Health and Medicine', write a letter in not more than 150 words.

- infrastructure unutilized
- pedestrians prone to accidents
- causes of apathy –
 - (a) accumulation of filth and garbage
 - (b) stagnant water
- need improvement in regular cleanliness and proper supervision.

2. The following lines from the poem 'Reaching Out' in the Main Course Book set you thinking about the need to educate adults. You decided to get involved in the adult education programme. Taking ideas from the unit 'Education', along with your own ideas write a letter to the Editor of a newspaper in about 150 words, encouraging youngsters to take part in the programme.


".... despite our illiteracy.

We still exist

But we have to know

Why we should become literate."

3. With the advent of plastic, a new and powerful enemy of environment has been born. For convenience-crazy humanity, plastic is a great boon, but for the environment, it is a menace. Tones of plastic waste litter the streets, choke drains, and pollute the environment. Waste plastic in rubbish dumps and landfill sites is causing the death of many animals. Plastic debris floats in the sea and endangers sea birds.



Look at the given cartoon. Write a letter to the Editor of 'Bharat Times', protesting against the growing plastic menace and giving suggestions about dealing with plastic waste. Write the letter in not more than 150 words, using the information given, your own ideas, and ideas from the unit 'Environment' in your Main Course Book.